

2025 Chemistry

Higher - Paper 1

Question Paper Finalised Marking Instructions

© Scottish Qualifications Authority 2025

These marking instructions have been prepared by examination teams for use by SQA appointed markers when marking external course assessments.

The information in this document may be reproduced in support of SQA qualifications only on a non-commercial basis. If it is reproduced, SQA must be clearly acknowledged as the source. If it is to be reproduced for any other purpose, written permission must be obtained from permissions@sqa.org.uk.

Marking instructions for each question

Question	Answer	Mark
1.	А	1
2.	С	1
3.	В	1
4.	D	1
5.	В	1
6.	А	1
7.	В	1
8.	D	1
9.	С	1
10.	В	1
11.	D	1
12.	В	1
13.	С	1
14.	А	1
15.	А	1
16.	D	1
17.	D	1
18.	В	1
19.	D	1
20.	С	1
21.	D	1
22.	А	1
23.	С	1
24.	С	1
25.	A	1

[END OF MARKING INSTRUCTIONS]



2025 Chemistry

Higher - Paper 2

Question Paper Finalised Marking Instructions

© Scottish Qualifications Authority 2025

These marking instructions have been prepared by examination teams for use by SQA appointed markers when marking external course assessments.

The information in this document may be reproduced in support of SQA qualifications only on a non-commercial basis. If it is reproduced, SQA must be clearly acknowledged as the source. If it is to be reproduced for any other purpose, written permission must be obtained from permissions@sqa.org.uk.



General marking principles for Higher Chemistry

Always apply these general principles. Use them in conjunction with the detailed marking instructions, which identify the key features required in candidates' responses.

- (a) Always use positive marking. This means candidates accumulate marks for the demonstration of relevant skills, knowledge and understanding; marks are not deducted for errors or omissions.
- (b) If a candidate response does not seem to be covered by either the principles or detailed marking instructions, and you are uncertain how to assess it, you must seek guidance from your team leader.
- (c) Do not award half marks.
- (d) Where a candidate makes an error at an early stage in a multi-stage calculation, award marks for correct follow-on working in subsequent stages, unless the error significantly reduces the complexity of the remaining stages. Apply the same principle for questions that require several stages of non-mathematical reasoning. The exception to this rule is where the marking instructions for a numerical question assign separate 'concept marks' and an 'arithmetic mark'. In such situations, the marking instructions will give clear guidance on the assignment or partial marks.
- (e) Unless a numerical question specifically requires evidence of working to be shown, award full marks for a correct final response (including units) on its own.
- (f) Candidates may fully access larger mark allocations whether their responses are in continuous prose, linked statements, or a series of developed bullet points.
- (g) Do not deduct marks for inaccurate or unconventional spelling or vocabulary as long as the meaning of the word(s) is conveyed. For example, responses that include 'distilling' for 'distillation', or 'it gets hotter' for 'the temperature rises', are acceptable.
- (h) In many questions, the unit in which the answer is to be expressed is given. In these questions, the candidate does not need to state a unit in their answer; but if they do, the unit must be correct. The full mark allocation cannot be awarded if an incorrect unit is shown. In these questions, incorrect units would only be penalised once in any paper.
- (i) If a correct response is followed by a wrong response, award no marks. For example in response to the question, 'State the colour seen when blue Fehling's solution is warmed with an aldehyde', do not award marks for the response 'red green'. However, if a correct response is followed by additional information which does not conflict with that, ignore the additional information, whether correct or not. For example in response to a question concerned with melting point, 'State why the tube should not be made of copper', the response 'Copper has a low melting point and is coloured grey' would gain marks.
- (j) Award full marks for the correct response to a calculation without working. Award partial marks, as shown in the detailed marking instructions, when working is given but the final response is incorrect. An exception is when candidates are asked to 'Find, by calculation' do not award full marks for the correct response without working.
- (k) Ignore the omission of one H atom from a full structural formula provided the bond is shown.
- (I) Award marks for a symbol or correct formula in place of a name unless stated otherwise in the detailed marking instructions.

- (m) When formulae of ionic compounds are given as responses, candidates only need to show ion charges if these have been specifically asked for. However, if ion charges are shown, they must be correct. If incorrect charges are shown, do not award marks.
- (n) If an answer comes directly from the text of the question, do not award marks. For example, in response to the question, 'A student found that 0.05 mol of propane, C_3H_8 burned to give 82.4 kJ of energy. $C_3H_8(g) + 5O_2(g) \rightarrow 3CO_2(g) + 4H_2O(\ell)$. Name the kind of enthalpy change that the student measured', do not award marks for 'burning' since the word 'burned' appears in the text.
- (o) A guiding principle in marking is to give credit for correct elements of a response rather than to look for reasons not to give marks.

Example 1: The structure of a hydrocarbon found in petrol is shown below.

$$CH_3$$
 $CH_3-CH_2-CH-CH_2-CH_3-CH_3$

Name the hydrocarbon

• Award the full mark for '3, methyl-hexane', although the punctuation is not correct.

Example 2: A student measured the pH of four carboxylic acids to find out how their strength is related to the number of chlorine atoms in the molecule. The results are shown.

CH₃COOH	1.65
CH₂ClCOOH	1.27
CHCl₂COOH	0.90
CCl₃COOH	0.51

Describe the relationship between the number of chlorine atoms in the molecule and the strengths of the acids.

- Award the full mark for a response such as 'the more Cl₂, the stronger the acid', although not completely correct.
- (p) Unless the question is clearly about a non-chemistry issue, for example costs in an industrial chemical process, do not award marks for a non-chemical response. For example, in response to the question, 'Why does the (catalytic) converter have a honeycomb structure?', do not award a mark for 'To make it work'. This response may be correct but it is not a chemical response.

- (q) Only award marks for a valid response to the question asked. Where candidates are asked to:
 - identify, name, give or state, they must only name or present in brief form.
 - **describe**, they must provide a statement or structure of characteristics and/or features.
 - **explain**, they must relate cause and effect and/or make relationships between things clear.
 - **compare**, they must demonstrate knowledge and understanding of the similarities and/or differences between things.
 - complete, they must finish a chemical equation or fill in a table with information.
 - **determine** or **calculate**, they must determine a number from given facts, figures or information.
 - draw, they must draw a diagram or structural formula, for example 'Draw a diagram to show the part of a poly(propene) molecule formed from two propene molecules.'
 - estimate, they must determine an approximate value for something.
 - predict, they must suggest what may happen based on available information.
 - evaluate, they must make a judgement based on criteria.
 - **suggest**, they must apply their knowledge and understanding of chemistry to a new situation. A number of responses are acceptable: award marks for any suggestions that are supported by knowledge and understanding of chemistry.
 - use their knowledge of chemistry or aspect of chemistry to comment on, they must apply their skills, knowledge and understanding to respond appropriately to the problem/situation presented (for example by making a statement of principle(s) involved and/or a relationship or equation, and applying these to respond to the problem/situation). Candidates gain marks for the breadth and/or depth of their conceptual understanding.
 - write, they must complete a chemical or word equation, for example 'Write the word equation for the complete combustion of ethanol.'

Page 6

Marking instructions for each question

C	Question		Expected response	Max mark	Additional guidance
1.	(a)	(i)	$Na(g) \longrightarrow Na^+(g) + e^-$	1	State symbols must be shown, negative sign on electron not required.
		(ii)	1735 (kJ mol ⁻¹)	1	No units required. No mark can be awarded for correct answer if wrong unit is given (where no unit required, wrong units would only be penalised once in any paper). kJ is acceptable in place of kJ mol-1 (KJ or Kj or KJ mol-1 or Kj mol-1 accepted).
		(iii)	The second ionisation energy (in sodium) involves removal of an electron from an (electron) shell that is inner/full/(more) stable/closer to the nucleus.	2	Correct statements made about the 2 nd ionisation energy/electron of magnesium can also be credited as long as magnesium is named.
			OR Second electron is removed from an (electron) shell that is inner/full/(more) stable/closer to the nucleus. (1 mark) The second electron is less screened/the second (electron) shell is less screened. OR The second electron is more strongly attracted to/pulled towards the nucleus. (1 mark)		Stating that the 2 nd electron requires more energy than the 1 st electron is not sufficient on its own. Shielding is acceptable in place of screening. Energy level is acceptable in place of electron shell. Increased attraction of the electron for the nucleus would be considered cancelling.
		(iv)	(They have the same number of occupied electron shells but) there are more protons/increased nuclear charge of the Mg ²⁺ ion.	1	

Looking for more resources? Visit https://sqa.my/ - Scotland's #1 Past Paper Archive

Q	Question		Expected response	Max mark	Additional guidance
1.	(b)	(i)	(Ionic bonds are) the electrostatic attraction between positive and negative ions. OR (An ionic bond is) the electrostatic attraction between oppositely charged ions.	1	Accept reference to a single ionic bond being the electrostatic attraction between a positive ion and a negative ion.
		(ii)	6 bananas (if working shown must be correct) Partial marks Mass of potassium in 7 day dose= 8 × 7 × 0.0012 × 39.1 = 2.62752 (g) (1 mark) Any calculated mass of potassium divided by 0.450. (1 mark)	2	2627.52 (mg) accepted for partial mark.

Q	Question		Expected re	esponse	Max mark	Additional guidance
2.	(a)		Bonding		2	Covalent lattice is accepted for (covalent) network.
			metallic			(covatent) network.
			covalent			
			covalent			
			covalent			
				(1 mark)		
			Structure			
			lattice			
			(covalent) networ	k		
			molecular			
			(discrete) molecular/molecul	le		
				(1 mark)		
	(b)	(i)	(The covalent radius)) decreases.	1	
		(ii)	Argon does not form bonds.	(covalent)	1	Argon is unreactive/monatomic is not accepted on its own.
	(c)		(van der Waals) force stronger (going down	_	3	Mention of increasing/stronger covalent/ionic/metallic bonds - cancels the first mark.
			LDFs are the forces (the atoms).	broken between (1 mark)		This mark can only be awarded if no other forces are mentioned (as being broken).
			The number of election (and so the stronger			Mention of (breaking) bonds/forces between molecules cancels one mark.

Qı	uestion	Expected response	Max mark	Additional guidance
3.	Jestion	Award 1 mark where the candidate has demonstrated, at an appropriate level, a limited understanding of the chemistry involved. They have made some statement(s) which are relevant to the situation, showing that they have understood at least a little of the chemistry within the problem. Award 2 marks where the candidate has demonstrated, at an appropriate level, a reasonable understanding of the chemistry involved. They make some statement(s) which are relevant to the situation, showing that they have understood the problem. Award 3 marks where the candidate has demonstrated, at an appropriate level, a good understanding of the chemistry involved. They show a good comprehension of the	mark 3	Additional guidance
		chemistry of the situation and provide a logically correct answer to the question posed. This type of response might include a statement of the principles involved, a relationship or an equation, and the application of these to respond to the problem. The answer does not need to be 'excellent' or 'complete' for the candidate to gain full marks. Award 0 marks where the candidate has not demonstrated, at an appropriate level, an understanding of the chemistry involved. There is no evidence that they have recognised the area of chemistry involved, or they have not given any statement of a relevant chemistry principle. Award this mark also if the candidate merely restates the chemistry given in the question.		

C)uesti	on	Expected response	Max mark	Additional guidance
4.	(a)	(i)	$C_3H_7OH + 4\frac{1}{2} O_2 \rightarrow 3CO_2 + 4H_2O$	1	Multiples not accepted.
		(ii)	-1050.04 (kJ mol ⁻¹)	3	1050.04 (kJ mol ⁻¹) (2 marks)
			If incorrect answer given, a maximum of two partial marks can be awarded for demonstration of the two concepts:		No units required. if wrong unit is given (where no unit required, wrong units would only be penalised once in any paper).
			1 mark for a demonstration of the correct use of the relationship Eh = cm∆T as shown by (4.18 × (an order of magnitude 150) × 13.9). (ignore units for this mark).		kJ is acceptable in place of kJ mol ⁻¹ (KJ or Kj or KJ mol ⁻¹ or Kj mol ⁻¹ accepted).
			1 mark for evidence of scaling up of a calculated value of energy released to one mole. This can be demonstrated by:		
			 correct scaling of calculated Eh from 0.498 to 60 g 0.498 g ← Eh 60 g 60 x Eh 0.498 0.498 (0.0083) ← Eh 		
			60 1 ←→ X		
		(iii)	 balance (alcohol) burner (copper) can/beaker thermometer measuring cylinder/volume measurement clamp stand/tripod/can suspension method Any four for 2 marks.	2	Bomb calorimeter worth maximum of 1 mark.
			Any two for 1 mark.		
		(iv)	Heat/energy loss to the surroundings OR Incomplete combustion. OR Evaporation of alcohol/fuel.	1	Evaporation on its own is not sufficient.
	(b)	(i)	1:3 OR 2:6	1	Hydrogen:oxygen ratio accepted if labelled.

C	Question		Expected response	Max mark	Additional guidance
4.	(b)	(ii) (A)	Acidified (potassium/sodium) dichromate.	1	Copper(II)oxide without mention of heating is insufficient for the mark.
			OR		
			Hot copper(II)oxide.		
		(ii) (B)	H H H O 	1	Correct structure of 2-methylpropanoic acid is also accepted.
		(ii) (C)	Butanone/butan-2-one.	1	Structure not accepted.
	(c)	(i)	Volumetric/standard flask.	1	
		(ii) (A)	Rinse with sodium hydroxide (before filling).	1	
		(ii) (B)	20.4 (cm ³)	1	Units are not required but must be correct if given. (wrong units would only be penalised once in any paper).

Qı	uesti	on	Expected response	Max mark	Additional guidance
4.	(c)	(ii) (C)	If incorrect answer given, partial marks can be awarded using a scheme of two "concept" marks, and one "arithmetic" mark. 1 mark for knowledge of the relationship between moles, concentration and volume. This could be shown by one of the following steps: Calculation of moles sodium hydroxide solution e.g. 0·105 x 0.0198 = 2.079 x 10 ⁻³ moles. OR Calculation of concentration of ethanoic acid e.g. 2.079 x 10 ⁻³ ÷ 0.025 OR Insertion of correct pairings of values for concentration and volume in a valid titration formula If the relationship between moles, concentration and volume is used more than once, it must be used correctly every time. 1 mark for knowledge of relationship between moles of sodium hydroxide and ethanoic acid. This could be shown by one of the following steps: Calculation of moles ethanoic acid from moles sodium hydroxide i.e. calculated moles of sodium hydroxide used again to calculate concentration of ethanoic acid. OR Insertion of correct stoichiometric values in a valid titration formula. 1 mark is awarded for correct arithmetic through the calculation. This mark can only be awarded if both concept marks have been awarded.	3	Units are not required but must be correct if given. (wrong units would only be penalised once in any paper)

Ç	Question		Expected response	Max mark	Additional guidance
4.	(c)	(iii) (A)	CH₃COO⁻K⁺	1	K ⁺ CH₃COO ⁻ is also acceptable.
		(iii) (B)	5.4 (g)	1	Units are not required but must be correct if given. (wrong units would only be penalised once in any paper)

Q	uestic	on	Expected response	Max mark	Additional guidance
5.	(a)		(Sucrose) is polar due to its hydroxyl groups.	2	(-)OH accepted as hydroxyl.
			OR		
			(Sucrose) can form hydrogen bonds due to its hydroxyl groups. (1 mark)		"Like dissolves like" on its own is not sufficient for the second mark.
			An explanation which links solubility of the molecule (sucrose) to the polarity of water/hydrogen bonding of water. (1 mark)		sufficient for the second mark.
	(b)	(i)	Tyr-Pro-Phe-Pro-Gly-Pro-Iso.	1	
			OR		
			Iso-Pro-Gly-Pro-Phe-Pro-Tyr.		
		(ii)	Essential (amino acid)	1	
	(c)	(i)	Ester(s)	1	Triglyceride accepted
		(ii) (A)	Glycerol/propane-1,2,3-triol	1	propan-1,2,3-triol accepted
		(ii) (B)	Palmitic acid	1	
		(iii)	Addition	1	Reduction accepted
	(d)	(i)	Ethanal	1	
		(ii)	Carbonyl	1	
		(iii) (A)	Diterpenoid	1	
		(iii) (B)	A correctly circled isoprene unit.	1	Including the OH is not accepted.
	(e)		3	1	

Q	Question		Expected response	Max mark	Additional guidance
6.	(a)		$H_2O_2(aq) + 2H^+(aq) + 2e^- \rightarrow 2H_2O(\ell)$	1	Ignore state symbols. Charge on electron not required.
	(b)		$2S_2O_3^{2-}(aq) + I_2(aq) \rightarrow S_4O_6^{2-}(aq) + 2I^{-}(aq)$	1	Ignore state symbols.
	(c)	(i)	10	1	
		(ii)	To keep the concentration (of the reactants) constant.	1	
			OR		
			Adding water will change/affect/dilute/decrease the concentration (of the reactants).		
	(d)	(i)	(As concentration doubles the rate of reaction) doubles.	1	
		(ii)	16.129 (s), 16.13(s)	1	Units are not required but must be correct if given. (wrong units would only be penalised once in any paper)
		(iii)	The more particles/ions (in a given volume). (1 mark)	2	
			More successful collisions (and the faster the reaction). (1 mark)		

Q	Question		Expected response	Max mark	Additional guidance	
7.	(a)		1129.4 (kJ mol ⁻¹) Partial marks Correctly retrieving 3 bond enthalpies from data book (243, 804, 338). (1 mark) Correct manipulation of incorrect bond enthalpies (must include the given ΔH = -107.6 kJ mol ⁻¹). (1 mark)	2	Retrieval of a fourth bond enthalpy (ie 360 for CO) is not considered cancelling for the first mark. Units not required, but if stated must be correct. Incorrect units would only be penalised once per paper. kJ is acceptable in place of kJ mol ⁻¹ (KJ or Kj or KJ mol ⁻¹ or Kj mol ⁻¹ accepted).	
	(b)		(Cooling favours) exothermic reaction. (1 mark) Cooling below 100 $^{\circ}$ C removes $H_2O(g)$ from equilibrium as $H_2O(\ell)$ (1 mark)	2		
	(c)	(i)	 mark for workable apparatus showing reaction of hydrochloric acid with calcium carbonate. mark for showing gas being passed through sealed container of sodium hydroxide. 	2	Treat each mark and associated labels separately.	
		(ii)	16.92/16.9/17(%) Partial marks Correct use of atom economy relationship without correct use of stoichiometry (working must be shown). (1 mark) Correct working with no correct answer given. (1 mark) Incorrect use of stoichiometry. Answer and working must be shown. (1 mark) 0.1692 (1 mark)	2		

Q	Question		Expected response	Max mark	Additional guidance
8.	(a)	(i)	198.4 (g) Partial marks Calculating number of moles of SnO ₂ required (1.316). (1 mark) OR Calculating the theoretical mass of tin required to produce 100 g at 64% yield (156.25). (1 mark) OR Calculating mass of SnO ₂ required for 100% yield (126.96). (1 mark) OR Follow through from an incorrectly calculated 100% yield. (1 mark)	2	
		(ii)	Carbon is in excess by 0.083 (moles). Partial marks Moles of SnO ₂ = 0.167 AND moles of C = 0.25 (1 mark) OR Allow for follow through from correctly applying mole ratio and subtraction to calculated number of moles (1 mark) By proportion 25.2 → 2.007 grams (1 mark) OR Allow for follow through from calculated mass of excess carbon converted to moles. (1 mark)	2	A maximum of 1 mark can be awarded if carbon is not stated as being the reactant in excess. 1 mark can be awarded for identification of tin(IV) oxide as the reactant in excess if supported by candidates working.

8. (a			mark	Additional guidance
	(a) (ii	Partial marks Calculating moles of CO₂ (1 mark) Correctly applying mole ratio to work out a mass for an incorrect number of moles. (1 mark) By proportion 150.7 g ← 80 litres (1 mark)	2	
(t	(b) (i) (ii		1	

Question			Expected response	Max mark	Additional guidance
9.	(a)	(i)	Propagation	1	
		(ii)	An (unpaired) electron	1	Do not accept electrons.
	(b)		Denaturing	1	
	(c)		Cosmetics/food products/plastics	1	
	(d)	(i)	Oxygen/gas is escaping	1	
		(ii)	 Two from: mass/volume/surface area/size of potato discs concentration of H₂O₂ temperature (of H₂O₂ or pH solution) 	2	Either from • type/variety of potato, or • (discs from the) same potato Accepted for 1 mark Weight is accepted for mass (of potato discs).
		(iii)	As the pH increases the rate (of reaction) increases up to pH 10/ reaches a maximum/optimum at pH 10 (and then decreases). OR As the pH increases the rate (of reaction) increases and then decreases.	1	Rate must be mentioned.

Qı	Question		Expected response	Max mark	Additional guidance
10.	(a)	(i)	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	
		(ii)	Bromine solution would quickly/rapidly decolourise with CAP/capsaicin. OR Bromine solution will decolourise with CAP/capsaicin but not with DHC/dihydrocapsaicin. OR Bromine solution will not decolourise with DHC/dihydrocapsaicin but will with CAP/capsaicin.	1	
	(b)		One from:	1	An answer referring to a named capsaicinoid is accepted. It/they is taken as referring to capsaicinoid.
	(c)		135 (s)	1	2.25 minutes is not accepted Sec(s) is not accepted as unit.
	(d)		Partial marks Mass of CAP in cream = 0.015 (g). OR Evidence of a calculated mass of CAP × (£)1.93/193(p). OR 60 g of CAP costs £115.82 OR 0.025% of a calculated cost of 60 g CAP OR 60 x £48.26 1000	2	£0.03 is accepted. Rounding of the final answer to the nearest correct monetary unit.
	(e)		$C_6H_8O_6 \rightarrow C_6H_6O_6 + 2H^+ + 2e^-$	1	State symbols are not required, but if shown must be correct. Negative sign on electron not required.

Question	Expected response	Max mark	Additional guidance
11.	Award 1 mark where the candidate has demonstrated, at an appropriate level, a limited understanding of the chemistry involved. They have made some statement(s) which are relevant to the situation, showing that they have understood at least a little of the chemistry within the problem. Award 2 marks where the candidate has demonstrated, at an appropriate level, a reasonable understanding of the chemistry involved. They make some statement(s) which are relevant to the situation, showing that they have understood the problem.	3	
	Award 3 marks where the candidate has demonstrated, at an appropriate level, a good understanding of the chemistry involved. They show a good comprehension of the chemistry of the situation and provide a logically correct answer to the question posed. This type of response might include a statement of the principles involved, a relationship or an equation, and the application of these to respond to the problem. The answer does not need to be 'excellent' or 'complete' for the candidate to gain full marks. Award 0 marks where the candidate has not demonstrated, at an appropriate level, an understanding of the chemistry involved. There is no evidence that they have recognised the area of chemistry involved, or they have not given any statement of a relevant chemistry principle. Award this mark also if the candidate merely restates the chemistry given in the question.		

Question			Expected response	Max mark	Additional guidance
12.	(a)		2,7-dimethyloctane	1	
	(b)		H	1	Any correct structure for 4-methylhex-2-ene
	(c)		OR OH	1	

[END OF MARKING INSTRUCTIONS]