



National
Qualifications
2017

2017 Mathematics Paper 1 (Non-calculator)

Higher

Finalised Marking Instructions

© Scottish Qualifications Authority 2017

The information in this publication may be reproduced to support SQA qualifications only on a non-commercial basis. If it is reproduced, SQA should be clearly acknowledged as the source. If it is to be used for any other purpose, written permission must be obtained from permissions@sqa.org.uk.

Where the publication includes materials from sources other than SQA (secondary copyright), this material should only be reproduced for the purposes of examination or assessment. If it needs to be reproduced for any other purpose it is the centre's responsibility to obtain the necessary copyright clearance. SQA's NQ Assessment team may be able to direct you to the secondary sources.

These marking instructions have been prepared by examination teams for use by SQA appointed markers when marking external course assessments. This publication must not be reproduced for commercial or trade purposes.



General marking principles for Higher Mathematics

This information is provided to help you understand the general principles you must apply when marking candidate responses to questions in this Paper. These principles must be read in conjunction with the detailed marking instructions, which identify the key features required in candidate responses.

For each question the marking instructions are generally in two sections, namely Illustrative Scheme and Generic Scheme. The illustrative scheme covers methods which are commonly seen throughout the marking. The generic scheme indicates the rationale for which each mark is awarded. In general, markers should use the illustrative scheme and only use the generic scheme where a candidate has used a method not covered in the illustrative scheme.

- (a) Marks for each candidate response must always be assigned in line with these general marking principles and the detailed marking instructions for this assessment.
- (b) Marking should always be positive. This means that, for each candidate response, marks are accumulated for the demonstration of relevant skills, knowledge and understanding: they are not deducted from a maximum on the basis of errors or omissions.
- (c) If a specific candidate response does not seem to be covered by either the principles or detailed marking instructions, and you are uncertain how to assess it, you must seek guidance from your Team Leader.
- (d) Credit must be assigned in accordance with the specific assessment guidelines.
- (e) One mark is available for each •. There are no half marks.
- (f) Working subsequent to an error must be **followed through**, with possible credit for the subsequent working, provided that the level of difficulty involved is approximately similar. Where, subsequent to an error, the working for a follow through mark has been eased, the follow through mark cannot be awarded.
- (g) As indicated on the front of the question paper, full credit should only be given where the solution contains appropriate working. Unless specifically mentioned in the marking instructions, a correct answer with no working receives no credit.
- (h) Candidates may use any mathematically correct method to answer questions except in cases where a particular method is specified or excluded.
- (i) As a consequence of an error perceived to be trivial, casual or insignificant, eg $6 \times 6 = 12$ candidates lose the opportunity of gaining a mark. However, note the second example in comment (j).
- (j)

- (j) Where a transcription error (paper to script or within script) occurs, the candidate should normally lose the opportunity to be awarded the next process mark, eg

This is a transcription error and so the mark is not awarded.		$x^2 + 5x + 7 = 9x + 4$
Eased as no longer a solution of a quadratic equation so mark is not awarded.		$x - 4x + 3 = 0$
Exceptionally this error is not treated as a transcription error as the candidate deals with the intended quadratic equation. The candidate has been given the benefit of the doubt and all marks awarded.		$x = 1$
		$x^2 + 5x + 7 = 9x + 4$
		$x - 4x + 3 = 0$
		$(x - 3)(x - 1) = 0$
		$x = 1 \text{ or } 3$

- (k) **Horizontal/vertical marking**

Where a question results in two pairs of solutions, this technique should be applied, but only if indicated in the detailed marking instructions for the question.

Example:

	• ⁵		• ⁶
• ⁵	$x = 2$	$x = -4$	
• ⁶	$y = 5$	$y = -7$	

Horizontal: • ⁵ $x = 2$ and $x = -4$	Vertical: • ⁵ $x = 2$ and $y = 5$
• ⁶ $y = 5$ and $y = -7$	• ⁶ $x = -4$ and $y = -7$

Markers should choose whichever method benefits the candidate, but **not** a combination of both.

- (l) In final answers, unless specifically mentioned in the detailed marking instructions, numerical values should be simplified as far as possible, eg:

$\frac{15}{12}$ must be simplified to $\frac{5}{4}$ or $1\frac{1}{4}$ $\frac{15}{0.3}$ must be simplified to 50 $\sqrt{64}$ must be simplified to 8*	$\frac{43}{1}$ must be simplified to 43 $\frac{4}{3}$ must be simplified to $\frac{4}{15}$
--	---

*The square root of perfect squares up to and including 100 must be known.

- (m) Commonly Observed Responses (COR) are shown in the marking instructions to help mark common and/or non-routine solutions. CORs may also be used as a guide when marking similar non-routine candidate responses.

(n) Unless specifically mentioned in the marking instructions, the following should not be penalised:

- Working subsequent to a correct answer
- Correct working in the wrong part of a question
- Legitimate variations in numerical answers/algebraic expressions, eg angles in degrees rounded to nearest degree
- Omission of units
- Bad form (bad form only becomes bad form if subsequent working is correct), eg $(x^3 + 2x^2 + 3x + 2)(2x + 1)$ written as $(x^3 + 2x^2 + 3x + 2) \times 2x + 1$

$2x^4 + 4x^3 + 6x^2 + 4x + x^3 + 2x^2 + 3x + 2$ written as $2x^4 + 5x^3 + 8x^2 + 7x + 2$ gains full credit

- Repeated error within a question, but not between questions or papers
- (o) In any 'Show that...' question, where the candidate has to arrive at a required result, the last mark of that part is not available as a follow-through from a previous error unless specified in the detailed marking instructions.
- (p) All working should be carefully checked, even where a fundamental misunderstanding is apparent early in the candidate's response. Marks may still be available later in the question so reference must be made continually to the marking instructions. The appearance of the correct answer does not necessarily indicate that the candidate has gained all the available marks.
- (q) Scored-out working which has not been replaced should be marked where still legible. However, if the scored out working has been replaced, only the work which has not been scored out should be marked.
- (r) Where a candidate has made multiple attempts using the same strategy and not identified their final answer, mark all attempts and award the lowest mark.

Where a candidate has tried different valid strategies, apply the above ruling to attempts within each strategy and then award the highest resultant mark.

For example:

Strategy 1 attempt 1 is worth 3 marks.	Strategy 2 attempt 1 is worth 1 mark.
Strategy 1 attempt 2 is worth 4 marks.	Strategy 2 attempt 2 is worth 5 marks.
From the attempts using strategy 1, the resultant mark would be 3.	From the attempts using strategy 2, the resultant mark would be 1.

In this case, award 3 marks.

Specific marking instructions for each question

Question			Generic scheme	Illustrative scheme	Max mark
1.	(a)		• ¹ evaluate expression	• ¹ 10	1
Notes:					
Commonly Observed Responses:					

Question			Generic scheme	Illustrative scheme	Max mark
1.	(b)		• ² interpret notation • ³ state expression for $g(f(x))$	• ² $g(5x)$ • ³ $2\cos 5x$	2
Notes:					
1. For $2\cos 5x$ without working, award both • ² and • ³ . 2. Candidates who interpret the composite function as either $g(x) \times f(x)$ or $g(x) + f(x)$ do not gain any marks. 3. $g(f(x)) = 10\cos x$ award • ² . However, $10\cos x$ with no working does not gain any marks. 4. $g(f(x))$ leading to $2\cos(5x)$ followed by incorrect 'simplification' of the function award • ² and • ³ .					
Commonly Observed Responses:					
Candidate A					
$g(f(x)) = 2\cos(5x)$ $= 10\cos(x)$			• ² ✓ • ³ ✓		

Question			Generic scheme	Illustrative scheme	Max mark
2.			<ul style="list-style-type: none"> •¹ state coordinates of centre •² find gradient of radius •³ state perpendicular gradient •⁴ determine equation of tangent 	<ul style="list-style-type: none"> •¹ (4, 3) •² $\frac{1}{3}$ •³ -3 •⁴ $y = -3x - 5$ 	4

Notes:

1. Accept $\frac{2}{6}$ for •².
2. The perpendicular gradient must be simplified at •³ or •⁴ stage for •³ to be available.
3. •⁴ is only available as a consequence of trying to find and use a perpendicular gradient.
4. At •⁴, accept $y + 3x + 5 = 0$, $y + 3x = -5$ or any other rearrangement of the equation where the constant terms have been simplified.

Commonly Observed Responses:

Question			Generic scheme	Illustrative scheme	Max mark
3.			<ul style="list-style-type: none"> •¹ start to differentiate •² complete differentiation 	<ul style="list-style-type: none"> •¹ $12(4x-1)^{11} \dots$ •² $\dots \times 4$ 	2

Notes:

1. •² is awarded for correct application of the chain rule.

Commonly Observed Responses:

<p>Candidate A</p> $\frac{dy}{dx} = 12(4x-1)^{11} \times 4 \quad \bullet^1 \checkmark \bullet^2 \checkmark$ $\frac{dy}{dx} = 36(4x-1)^{11}$ <p>Working subsequent to a correct answer: General Marking Principle (n)</p>	<p>Candidate B</p> $\frac{dy}{dx} = 36(4x-1)^{11} \quad \bullet^1 \times \bullet^2 \times$ <p>Incorrect answer with no working</p>
---	---

Question			Generic scheme	Illustrative scheme	Max mark
4.			<p>Method 1</p> <ul style="list-style-type: none"> •¹ use the discriminant •² apply condition and simplify •³ determine the value of k 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $4^2 - 4 \times 1 \times (k - 5)$ •² $36 - 4k = 0$ or $36 = 4k$ •³ $k = 9$ 	3
			<p>Method 2</p> <ul style="list-style-type: none"> •¹ communicate and express in factorised form •² expand and compare •³ determine the value of k 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ equal roots $\Rightarrow x^2 + 4x + (k - 5) = (x + 2)^2$ •² $x^2 + 4x + 4$ leading to $k - 5 = 4$ •³ $k = 9$ 	

Notes:

- At the •¹ stage, treat $4^2 - 4 \times 1 \times k - 5$ as bad form only if the candidate treats ' $k - 5$ ' as if it is bracketed in their next line of working. See Candidates A and B.
- In Method 1 if candidates use any condition other than 'discriminant = 0' then •² is lost and •³ is unavailable.

Commonly Observed Responses:

Candidate A		Candidate B	
$4^2 - 4 \times 1 \times k - 5$	• ¹ ✓	$4^2 - 4 \times 1 \times k - 5$	• ¹ ✗
$36 - 4k = 0$	• ² ✓	$11 - 4k = 0$	• ² ✓1
$k = 9$	• ³ ✓	$k = \frac{11}{4}$	• ³ ✓1

Question	Generic scheme	Illustrative scheme	Max mark
5. (a)	• ¹ evaluate scalar product	• ¹ 1	1

Notes:

Commonly Observed Responses:

Question	Generic scheme	Illustrative scheme	Max mark
5. (b)	• ² calculate $ \mathbf{u} $ • ³ use scalar product • ⁴ evaluate $\mathbf{u} \cdot \mathbf{w}$	• ² $\sqrt{27}$ • ³ $\sqrt{27} \times \sqrt{3} \times \cos \frac{\pi}{3}$ • ⁴ $\frac{9}{2}$ or 4.5	3

Notes:

1. Candidates who treat negative signs with a lack of rigour and arrive at $\sqrt{27}$ gain •².
2. Surds must be fully simplified for •⁴ to be awarded.

Commonly Observed Responses:

Question			Generic scheme	Illustrative scheme	Max mark
6.			Method 1 • ¹ equate composite function to x • ² write $h(h^{-1}(x))$ in terms of $h^{-1}(x)$ • ³ state inverse function	Method 1 • ¹ $h(h^{-1}(x)) = x$ • ² $(h^{-1}(x))^3 + 7 = x$ • ³ $h^{-1}(x) = \sqrt[3]{x-7}$ or $h^{-1}(x) = (x-7)^{\frac{1}{3}}$	3
			Method 2 • ¹ write as $y = x^3 + 7$ and start to rearrange • ² complete rearrangement • ³ state inverse function	Method 2 • ¹ $y - 7 = x^3$ • ² $x = \sqrt[3]{y-7}$ • ³ $h^{-1}(x) = \sqrt[3]{x-7}$ or $h^{-1}(x) = (x-7)^{\frac{1}{3}}$	3
			Method 3 • ¹ interchange variables • ² complete rearrangement • ³ state inverse function	Method 3 • ¹ $x = y^3 + 7$ • ² $y = \sqrt[3]{x-7}$ • ³ $h^{-1}(x) = \sqrt[3]{x-7}$ or $h^{-1}(x) = (x-7)^{\frac{1}{3}}$	3
Notes:					
1. $y = \sqrt[3]{x-7}$ (or $y = (x-7)^{\frac{1}{3}}$) does not gain • ³ .					
2. At • ³ stage, accept h^{-1} expressed in terms of any dummy variable eg $h^{-1}(y) = \sqrt[3]{y-7}$.					
3. $h^{-1}(x) = \sqrt[3]{x-7}$ or $h^{-1}(x) = (x-7)^{\frac{1}{3}}$ with no working gains 3/3.					

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
Candidate A			
$x \rightarrow x^3 \rightarrow x^3 + 7 = h(x)$ $\wedge 3 \rightarrow +7$ $\therefore -7 \rightarrow \sqrt[3]{}$ $\sqrt[3]{x-7}$ $h^{-1}(x) = \sqrt[3]{x-7}$		<p>•¹✓ awarded for knowing to perform the inverse operations in reverse order</p> <p>•²✓</p> <p>•³✓</p>	
Candidate B - BEWARE		Candidate C	
$h'(x) = \dots$ • ³ ✗		$h^{-1}(x) = \sqrt[3]{x} - 7$ • ³ ✗ With no working 0/3	

Question			Generic scheme	Illustrative scheme	Max mark
7.			<ul style="list-style-type: none"> •¹ find midpoint of AB •² demonstrate the line is vertical •³ state equation 	<ul style="list-style-type: none"> •¹ (2,7) •² m_{median} undefined •³ $x = 2$ 	3

Notes:

1. $m_{\text{median}} = \frac{\pm 4}{0}$ alone is not sufficient to gain •². Candidates must use either 'vertical' or 'undefined'. However •³ is still available.
2. ' $m_{\text{median}} = \frac{4}{0} \times$ ', ' $m_{\text{median}} = \frac{4}{0}$ impossible' ' $m_{\text{median}} = \frac{4}{0}$ infinite' are **not** acceptable for •².
However, if these are followed by either 'vertical' or 'undefined' then award •², and •³ is still available.
3. ' $m_{\text{median}} = \frac{4}{0} = 0$ undefined' ' $m_{\text{median}} = \frac{4}{0}$ undefined' are **not** acceptable for •².
4. •³ is not available as a consequence of using a numeric gradient; however, see notes 5 and 6.
5. For candidates who find an incorrect midpoint (a,b) , using the coordinates of A and B and find the 'median' through C without any further errors award 1/3. However, if $a = 2$, then both •² and •³ are available.
6. For candidates who find $15y = 2x + 121$ (median through B) or $3y = 2x + 21$ (median through A) award 1/3.

Commonly Observed Responses:

Candidate A $(2,7)$ • ¹ ✓ $m = \frac{4}{0}$ $= 0$ undefined • ² ✗ $x = 2$ • ³ ✓1	Candidate B $(2,7)$ • ¹ ✓ $m = \frac{4}{0}$ $= 0$ • ² ✗ $y = 7$ • ³ ✓2	Candidate C $(2,7)$ • ¹ ✓ $m = \frac{4}{0}$ • ² ^ $y - 7 = \frac{4}{0}(x - 2)$ $0 = 4x - 8$ $x = 2$ • ³ ✗
Candidate D $(2,7)$ • ¹ ✓ Median passes through $(2,7)$ and $(2,11)$ • ² ✗ $x = 2$ • ³ ✓1	Candidate E $(2,7)$ • ¹ ✓ Both coordinates have an x value 2 \Rightarrow vertical line $x = 2$ • ² ✓ • ³ ✓	

Question			Generic scheme	Illustrative scheme	Max mark
8.			<p>•¹ write in differentiable form</p> <p>•² differentiate</p> <p>•³ evaluate derivative</p>	<p>•¹ $\frac{1}{2}t^{-1}$</p> <p>•² $-\frac{1}{2}t^{-2}$</p> <p>•³ $-\frac{1}{50}$</p>	3

Notes:

1. Candidates who arrive at an expression containing more than one term at •¹ award 0/3.
2. •² is only available for differentiating a term containing a negative power of t .

Commonly Observed Responses:

Candidate A		Candidate B		Candidate C	
$2t^{-1}$	• ¹ ✗	$2t^{-1}$	• ¹ ✗	$-\frac{1}{2}t^{-2}$	• ¹ ✓ implied by • ² ✓
$-2t^{-2}$	• ² ✓1	$-2t^{-2}$	• ² ✓1	$-\frac{1}{50}$	• ³ ✓
$-\frac{2}{25}$	• ³ ✓1	$-\frac{1}{50}$	• ³ ✗		
Candidate D		Candidate E		Candidate F	
				Bad form of chain rule	
$(2t)^{-1}$	• ¹ ✓	$(2t)^{-1}$	• ¹ ✓	$2t^{-1}$	• ¹ ✓
$-(2t)^{-2}$	• ² ✗	$-(2t)^{-2}$	• ² ✗	$\underline{\underline{-2t^{-2}}} \times 2$	• ² ✓
$-\frac{1}{100}$	• ³ ✓1	$-\frac{2}{25}$	• ³ ✗	$-\frac{1}{50}$	• ³ ✓
				Candidate G	
				$2t^{-1}$	• ¹ ✗
				$\underline{\underline{-2t^{-2}}} \times 2$	• ² ✗
				$-\frac{4}{25}$	• ³ ✓1

Question			Generic scheme	Illustrative scheme	Max mark
9.	(a)		<ul style="list-style-type: none"> •¹ interpret information •² state the value of m 	<ul style="list-style-type: none"> •¹ $13 = 28m + 6$ stated explicitly or in a rearranged form •² $m = \frac{1}{4}$ or $m = 0.25$ 	2
Notes:					
1. Stating ' $m = \frac{1}{4}$ ', or simply writing ' $\frac{1}{4}$ ', with no other working gains only • ² .					
Commonly Observed Responses:					
Candidate A			Candidate B		
$13 = 28u_n + 6$			$28 = 13m + 6$		
$u_n = \frac{1}{4}$			$m = \frac{22}{13}$		
<ul style="list-style-type: none"> •¹ ✗ •² ✓1 			<ul style="list-style-type: none"> •¹ ✗ •² ✓1 		

Question			Generic scheme	Illustrative scheme	Max mark
9.	(b)	(i)	• ³ communicate condition for limit to exist	• ³ a limit exists as the recurrence relation is linear and $-1 < \frac{1}{4} < 1$	1

Notes:

2. For •³ accept:

any of $-1 < \frac{1}{4} < 1$ or $\left| \frac{1}{4} \right| < 1$ or $0 < \frac{1}{4} < 1$ with no further comment;

or statements such as:

“ $\frac{1}{4}$ lies between -1 and 1 ” or “ $\frac{1}{4}$ is a proper fraction”

3. •³ is not available for:

$-1 \leq \frac{1}{4} \leq 1$ or $\frac{1}{4} < 1$

or statements such as:

“It is between -1 and 1 .” or “ $\frac{1}{4}$ is a fraction.”

4. Candidates who state $-1 < m < 1$ can only gain •³ if it is **explicitly stated** that $m = \frac{1}{4}$ in part (a).

5. Do not accept ‘ $-1 < a < 1$ ’ for •³.

Commonly Observed Responses:

Candidate C			Candidate D		
(a)	$m = \frac{1}{4}$	• ¹ ✓ • ² ✓	(a)	$\frac{1}{4}$	• ¹ ✓ • ² ✓
(b)	$-1 < m < 1$	• ³ ✓	(b)	$-1 < m < 1$	• ³ ✗

Question			Generic scheme	Illustrative scheme	Max mark
9.	(b)	(ii)	<ul style="list-style-type: none"> •⁴ know how to calculate limit •⁵ calculate limit 	<ul style="list-style-type: none"> •⁴ $\frac{6}{1-\frac{1}{4}}$ or $L = \frac{1}{4}L + 6$ •⁵ 8 	2

Notes:

- Do not accept $L = \frac{b}{1-a}$ with no further working for •⁴.
- ⁴ and •⁵ are not available to candidates who conjecture that $L = 8$ following the calculation of further terms in the sequence.
- For $L = 8$ with no working, award 0/2.
- For candidates who use a value of m appearing ex nihilo or which is inconsistent with their answer in part (a) •⁴ and •⁵ are not available.

Commonly Observed Responses:

Candidate E - no valid limit

(a) $m = 4$ •¹ ✗

(b) $L = \frac{6}{1-4}$ •⁴ ✓1
 $L = -2$ •⁵ ✗

Question			Generic scheme	Illustrative scheme	Max mark
10.	(a)		<ul style="list-style-type: none"> •¹ know to integrate between appropriate limits •² use “upper - lower” •³ integrate •⁴ substitute limits •⁵ evaluate area 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $\int_0^2 \dots dx$ •² $\int_0^2 \left((x^3 - 4x^2 + 3x + 1) - (x^2 - 3x + 1) \right)$ •³ $\frac{x^4}{4} - \frac{5x^3}{3} + 3x^2$ •⁴ $\left(\frac{2^4}{4} - \frac{5 \times 2^3}{3} + 3 \times 2^2 \right) - (0)$ •⁵ $\frac{8}{3}$ 	
			<ul style="list-style-type: none"> •¹ know to integrate between appropriate limits for both integrals •² integrate both functions •³ substitute limits into both functions •⁴ evaluation of both functions •⁵ evidence of subtracting areas 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ $\int_0^2 \dots dx$ and $\int_0^2 \dots dx$ •² $\frac{x^4}{4} - \frac{4x^3}{3} + \frac{3x^2}{2} + x$ and $\frac{x^3}{3} - \frac{3x^2}{2} + x$ •³ $\left(\frac{2^4}{4} - \frac{4(2^3)}{3} + \frac{3(2^2)}{2} + 2 \right) - 0$ and $\left(\frac{2^3}{3} - \frac{3(2^2)}{2} + 2 \right) - 0$ •⁴ $\frac{4}{3}$ and $\frac{-4}{3}$ •⁵ $\frac{4}{3} - \frac{-4}{3} = \frac{8}{3}$ 	

5

Question	Generic scheme	Illustrative scheme	Max mark
Notes:			
<div>1. •¹ is not available to candidates who omit 'dx'.</div> <div>2. Treat the absence of brackets at •² stage as bad form only if the correct integral is obtained at •³. See Candidates A and B.</div> <div>3. Where a candidate differentiates one or more terms at •³, then •³, •⁴ and •⁵ are unavailable.</div> <div>4. Accept unsimplified expressions at •³ e.g. $\frac{x^4}{4} - \frac{4x^3}{3} + \frac{3x^2}{2} + x - \frac{x^3}{3} + \frac{3x^2}{2} - x$.</div> <div>5. Do not penalise the inclusion of '+c'.</div> <div>6. Candidates who substitute limits without integrating do not gain •³, •⁴ or •⁵.</div> <div>7. •⁴ is only available if there is evidence that the lower limit '0' has been considered.</div> <div>8. Do not penalise errors in substitution of $x=0$ at •³.</div>			
Commonly Observed Responses:			
<div>Candidate A</div> <div>•¹ ✓</div> <div>$\int_0^2 x^3 - 4x^2 + 3x + 1 - x^2 - 3x + 1 \, dx$</div> <div>$\frac{x^4}{4} - \frac{5x^3}{3} + 3x^2$</div> <div>•³ ✓ \Rightarrow •² ✓</div>		<div>Candidate B</div> <div>•¹ ✓</div> <div>$\int_0^2 x^3 - 4x^2 + 3x + 1 - x^2 - 3x + 1 \, dx$</div> <div>•² ✗</div> <div>$\frac{x^4}{4} - \frac{5x^3}{3} + 2x$</div> <div>•³ ✓1</div> <div>$\int \dots = -\frac{16}{3}$ cannot be negative so $= \frac{16}{3}$</div> <div>•⁵ ✗</div> <div>However, $\int \dots = -\frac{16}{3}$ so Area $= \frac{16}{3}$</div> <div>•⁵ ✓</div>	
Treating individual integrals as areas			
<div>Candidate C - Method 2</div> <div>•¹ ✓</div> <div>•² ✓</div> <div>•³ ✓</div> <div>$\frac{4}{3}$ and $-\frac{4}{3}$</div> <div>•⁴ ✓</div> <div>\therefore Area is $\frac{4}{3} - \left(-\frac{4}{3}\right) = \frac{8}{3}$</div> <div>•⁵ ✓</div>	<div>Candidate D - Method 2</div> <div>•¹ ✓</div> <div>•² ✓</div> <div>•³ ✓</div> <div>$\frac{4}{3}$ and $-\frac{4}{3}$</div> <div>•⁴ ✓</div> <div>$= \frac{4}{3}$</div> <div>\therefore Area is $\frac{4}{3} + \frac{4}{3} = \frac{8}{3}$</div> <div>•⁵ ✗</div>	<div>Candidate E - Method 2</div> <div>•¹ ✓</div> <div>•² ✓</div> <div>•³ ✓</div> <div>$\frac{4}{3}$ and $-\frac{4}{3}$</div> <div>•⁴ ✓</div> <div>Area cannot be negative</div> <div>\therefore Area is $\frac{4}{3} + \frac{4}{3} = \frac{8}{3}$</div> <div>•⁵ ✗</div>	

Question		Generic scheme	Illustrative scheme	Max mark
10.	(b)	<ul style="list-style-type: none"> •⁶ use “line - quadratic” •⁷ integrate •⁸ substitute limits and evaluate integral •⁹ state fraction 	<p>Method 1</p> <ul style="list-style-type: none"> •⁶ $\int \left((1-x) - (x^2 - 3x + 1) \right) dx$ •⁷ $-\frac{x^3}{3} + x^2$ •⁸ $\left(-\frac{2^3}{3} + 2^2 \right) - (0) = \frac{4}{3}$ •⁹ $\frac{1}{2}$ 	
		<ul style="list-style-type: none"> •⁶ use “cubic - line” •⁷ integrate •⁸ substitute limits and evaluate integral •⁹ state fraction 	<p>Method 2</p> <ul style="list-style-type: none"> •⁶ $\int \left((x^3 - 4x^2 + 3x + 1) - (1-x) \right) dx$ •⁷ $\frac{x^4}{4} - \frac{4x^3}{3} + 2x^2$ •⁸ $\left(\frac{2^4}{4} - 4 \times \frac{2^3}{3} + 2 \times 2^2 \right) - (0) = \frac{4}{3}$ •⁹ $\frac{1}{2}$ 	
		<ul style="list-style-type: none"> •⁶ integrate line •⁷ substitute limits and evaluate integral •⁸ evidence of subtracting integrals •⁹ state fraction 	<p>Method 3</p> <ul style="list-style-type: none"> •⁶ $\int (1-x) dx = \left[x - \frac{x^2}{2} \right]_0^2$ •⁷ $\left(2 - \frac{2^2}{2} \right) - (0) = 0$ •⁸ $0 - \left(-\frac{4}{3} \right) = \frac{4}{3}$ or $\frac{4}{3} - 0$ •⁹ $\frac{1}{2}$ 	

4

Question	Generic scheme	Illustrative scheme	Max mark
Notes:			
<p>IMPORTANT: Notes prefixed by *** may be subject to General Marking Principle (n). If a candidate has been penalised for the error in (a) then they must not be penalised a second time for the same error in (b).</p> <p>9. *** •⁶ is not available to candidates who omit 'dx'.</p> <p>10. In Methods 1 and 2 only, treat the absence of brackets at •⁶ stage as bad form only if the correct integral is obtained at •⁷.</p> <p>11. Candidates who have an incorrect expression to integrate at the •³ and •⁷ stage due solely to the absence of brackets lose •², but are awarded •⁶.</p> <p>12. Where a candidate differentiates one or more terms at •⁷, then •⁷, •⁸ and •⁹ are unavailable. *** In cases where Note 3 has applied in part (a), •⁷ is lost but •⁸ and •⁹ are available.</p> <p>13. In Methods 1 and 2 only, accept unsimplified expressions at •⁷ e.g. $x - \frac{x^2}{2} - \frac{x^3}{3} + \frac{3x^2}{2} - x$</p> <p>14. Do not penalise the inclusion of '+c'.</p> <p>15. *** •⁸ in Methods 1 and 2 and •⁷ in method 3 is only available if there is evidence that the lower limit '0' has been considered.</p> <p>16. At the •⁹ stage, the fraction must be consistent with the answers at •⁵ and •⁸ for •⁹ to be awarded.</p> <p>17. Do not penalise errors in substitution of $x = 0$ at •⁸ in Method 1 & 2 or •⁷ in Method 3.</p>			
Commonly Observed Responses:			

Question			Generic scheme	Illustrative scheme	Max mark
11.			<ul style="list-style-type: none"> •¹ determine the gradient of given line or of AB •² determine the other gradient •³ find a 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $\frac{2}{3}$ or $\frac{a-2}{12}$ •² $\frac{a-2}{12}$ or $\frac{2}{3}$ •³ 10 	3
			<ul style="list-style-type: none"> •¹ determine the gradient of given line •² equation of line and substitute •³ solve for a 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ $\frac{2}{3}$ stated or implied by •² •² $y-2 = \frac{2}{3}(x+7)$ $a-2 = \frac{2}{3}(5+7)$ •³ 10 	

Notes:

Commonly Observed Responses:

<p>Candidate A - using simultaneous equations</p> $m_{\text{line}} = \frac{2}{3} \quad \bullet^1 \checkmark$ $\left. \begin{array}{l} 3y = 2x + 20 \\ 3y = 2x - 10 + 3a \end{array} \right\} \bullet^2 \checkmark$ $0 = 0 + 30 - 3a$ $3a = 30$ $a = 10 \quad \bullet^3 \checkmark$	<p>Candidate B</p> $m_{AB} = \frac{a-2}{12} \quad \bullet^1 \checkmark$ $\frac{a-2}{12} = \underline{-2} \quad \bullet^2 \times$ $a = -22 \quad \bullet^3 \boxed{\checkmark 1}$	<p>Candidate C - Method 2</p> $\bullet^1 \checkmark$ $y-2 = \frac{2}{3}(x+7)$ $3y = 2x + 20$ $3y = 2 \times 5 + 20 \quad \bullet^2 \checkmark$ $3y = 30$ $y = 10$ <p>No mention of a $\bullet^3 \wedge$</p>
---	--	--

Question			Generic scheme	Illustrative scheme	Max mark
12.			<ul style="list-style-type: none"> •¹ use laws of logs •² write in exponential form •³ solve for a 	<ul style="list-style-type: none"> •¹ $\log_a 9$ •² $a^{\frac{1}{2}} = 9$ •³ 81 	3

Notes:

1. $\frac{36}{4}$ must be simplified at •¹ or •² stage for •¹ to be awarded.
2. Accept $\log 9$ at •¹.
3. •² may be implied by •³.

Commonly Observed Responses:

Candidate A	Candidate B	Candidate C
$\log_a 144$ • ¹ ✗ $a^{\frac{1}{2}} = 144$ • ² ✓1 $a = 12$ • ³ ✗	$\log_a 32$ • ¹ ✗ $a^{\frac{1}{2}} = 32$ • ² ✓1 $\bullet^3 \wedge$	$\log_a 9$ • ¹ ✓ $a = 9^{\frac{1}{2}}$ • ² ✗ $a = 3$ • ³ ✓2
Candidate D $2\log_a 36 - 2\log_a 4 = 1$ $\log_a 36^2 - \log_a 4^2 = 1$ • ¹ ✓ $\log_a \frac{36^2}{4^2} = 1$ $\log_a 81 = 1$ • ² ✓ $a = 81$ • ³ ✓		

Question			Generic scheme	Illustrative scheme	Max mark
13.			<p>•¹ write in integrable form</p> <p>•² start to integrate</p> <p>•³ process coefficient of x</p> <p>•⁴ complete integration and simplify</p>	<p>•¹ $(5-4x)^{-\frac{1}{2}}$</p> <p>•² $\frac{(5-4x)^{\frac{1}{2}}}{\frac{1}{2}} \dots$</p> <p>•³ $\dots \times \frac{1}{(-4)}$</p> <p>•⁴ $-\frac{1}{2}(5-4x)^{\frac{1}{2}} + c$</p>	4

Notes:

1. For candidates who differentiate throughout, only •¹ is available.
2. For candidates who 'integrate the denominator' without attempting to write in integrable form award 0/4.
3. If candidates start to integrate individual terms within the bracket or attempt to expand a bracket no further marks are available.
4. '+c' is required for •⁴.

Commonly Observed Responses:

<p>Candidate A</p> <p>$(5-4x)^{-\frac{1}{2}}$ •¹ ✓</p> <p>$\frac{(5-4x)^{\frac{1}{2}}}{\frac{1}{2}}$ •² ✓ •³ ^</p> <p>$2(5-4x)^{\frac{1}{2}} + c$ •⁴ ✓2</p>	<p>Candidate B</p> <p>$(5-4x)^{\frac{1}{2}}$ •¹ ✗</p> <p>$\frac{(5-4x)^{\frac{3}{2}}}{\frac{3}{2}} \times \frac{1}{(-4)}$ •² ✓1 •³ ✓</p> <p>$-\frac{(5-4x)^{\frac{3}{2}}}{6} + c$ •⁴ ✓1</p>
<p>Candidate C</p> <p>Differentiate in part:</p> <p>$(5-4x)^{-\frac{1}{2}}$ •¹ ✓</p> <p>$-\frac{1}{2}(5-4x)^{-\frac{3}{2}} \times \frac{1}{(-4)}$ •² ✗ •³ ✓</p> <p>$\frac{1}{8}(5-4x)^{-\frac{3}{2}} + c$ •⁴ ✓1</p>	<p>Candidate D</p> <p>Differentiate in part:</p> <p>$(5-4x)^{-\frac{1}{2}}$ •¹ ✓</p> <p>$\frac{(5-4x)^{\frac{1}{2}}}{\frac{1}{2}} \times (-4)$ •² ✓ •³ ✗</p> <p>$-8(5-4x)^{\frac{1}{2}} + c$ •⁴ ✓1</p>

Question			Generic Scheme	Illustrative Scheme	Max Mark
14.	(a)		<ul style="list-style-type: none"> •¹ use compound angle formula •² compare coefficients •³ process for k •⁴ process for a and express in required form 	<ul style="list-style-type: none"> •¹ $k \sin x^\circ \cos a^\circ - k \cos x^\circ \sin a^\circ$ stated explicitly •² $k \cos a^\circ = \sqrt{3}, k \sin a^\circ = 1$ stated explicitly •³ $k = 2$ •⁴ $2\sin(x - 30)^\circ$ 	4

Notes:

- Accept $k(\sin x^\circ \cos a^\circ - \cos x^\circ \sin a^\circ)$ for •¹. Treat $k \sin x^\circ \cos a^\circ - \cos x^\circ \sin a^\circ$ as bad form only if the equations at the •² stage both contain k .
- Do not penalise the omission of degree signs.
- $2\sin x^\circ \cos a^\circ - 2\cos x^\circ \sin a^\circ$ or $2(\sin x^\circ \cos a^\circ - \cos x^\circ \sin a^\circ)$ is acceptable for •¹ and •³.
- In the calculation of $k = 2$, do not penalise the appearance of -1 .
- Accept $k \cos a^\circ = \sqrt{3}, -k \sin a^\circ = -1$ for •².
- ² is not available for $k \cos x^\circ = \sqrt{3}, k \sin x^\circ = 1$, however, •⁴ is still available.
- ³ is only available for a single value of $k, k > 0$.
- ³ is not available to candidates who work with $\sqrt{4}$ throughout parts (a) and (b) without simplifying at any stage.
- ⁴ is not available for a value of a given in radians.
- Candidates may use any form of the wave equation for •¹, •² and •³, however, •⁴ is only available if the value of a is interpreted in the form $k \sin(x - a)^\circ$
- Evidence for •⁴ may only appear as a label on the graph in part (b).

Commonly Observed Responses:

Responses with missing information in working:

Candidate A		Candidate B	
$2\cos a = \sqrt{3}$ $2\sin a = 1$ $\tan a = \frac{1}{\sqrt{3}}, a = 30$ $2\sin(x - 30)^\circ$	<ul style="list-style-type: none"> •¹ ^ •² ✓ •³ ✓ •⁴ ✓ 	$k \sin x \cos a - k \cos x \sin a$ • ¹ ✓ $\cos a = \sqrt{3}$ $\sin a = 1$ • ² ✗ $\tan a = \frac{1}{\sqrt{3}}$ $a = 30$ <div style="border: 1px solid black; padding: 2px; display: inline-block;"> Not consistent with equations at •². </div> $2\sin(x - 30)^\circ$ • ³ ✓ • ⁴ ✗	

Question	Generic Scheme	Illustrative Scheme	Max Mark
Responses with the correct expansion of $k \sin(x-a)^\circ$ but errors for either \bullet^2 or \bullet^4 .			
Candidate C $k \cos a = \sqrt{3}, k \sin a = 1 \quad \bullet^2 \checkmark$ $\tan a = \sqrt{3} \quad \bullet^4 \times$ $a = 60$	Candidate D $k \cos a = 1, k \sin a = \sqrt{3} \quad \bullet^2 \times$ $\tan a = \sqrt{3}$ $a = 60$ $2 \sin(x-60)^\circ \quad \bullet^4 \boxed{\checkmark 1}$	Candidate E $k \cos a = \sqrt{3}, k \sin a = -1 \quad \bullet^2 \times$ $\tan a = -\frac{1}{\sqrt{3}}, a = 330$ $2 \sin(x-330)^\circ \quad \bullet^4 \boxed{\checkmark 1}$	
Responses with the incorrect labelling; $k \sin A \cos B - k \cos A \sin B$ from formula list.			
Candidate F $k \sin A \cos B - k \cos A \sin B \quad \bullet^1 \times$ $k \cos a = \sqrt{3}$ $k \sin a = 1 \quad \bullet^2 \checkmark$ $\tan a = \frac{1}{\sqrt{3}}, a = 30$ $2 \sin(x-30)^\circ \quad \bullet^3 \checkmark \bullet^4 \checkmark$	Candidate G $k \sin A \cos B - k \cos A \sin B \quad \bullet^1 \times$ $k \cos x = \sqrt{3}$ $k \sin x = 1 \quad \bullet^2 \times$ $\tan x = \frac{1}{\sqrt{3}}, x = 30$ $2 \sin(x-30)^\circ \quad \bullet^3 \checkmark \bullet^4 \boxed{\checkmark 1}$	Candidate H $k \sin A \cos B - k \cos A \sin B \quad \bullet^1 \times$ $k \cos B = \sqrt{3}$ $k \sin B = 1 \quad \bullet^2 \times$ $\tan B = \frac{1}{\sqrt{3}}, B = 30$ $2 \sin(x-30)^\circ \quad \bullet^3 \checkmark \bullet^4 \boxed{\checkmark 1}$	

Question			Generic scheme	Illustrative scheme	Max mark
14.	(b)		<ul style="list-style-type: none"> •⁵ roots identifiable from graph •⁶ coordinates of both turning points identifiable from graph •⁷ y-intercept and value of y at $x = 360$ identifiable from graph 	<ul style="list-style-type: none"> •⁵ 30 and 210 •⁶ (120, 2) and (300, -2) •⁷ -1 	3
Notes:					
<p>12. •⁵, •⁶ and •⁷ are only available for attempting to draw a “sine” graph with a period of 360°.</p> <p>13. Ignore any part of a graph drawn outwith $0 \leq x \leq 360$.</p> <p>14. Vertical marking is not applicable to •⁵ and •⁶.</p> <p>15. Candidates sketch arrived at in (b) must be consistent with the equation obtained in (a), see also candidates I and J.</p> <p>16. For any incorrect horizontal translation of the graph of the wave function arrived at in part(a) only •⁶ is available.</p>					
Commonly Observed Responses:					
Candidate I			Candidate J		
(a) $2\sin(x - 30)$ correct equation			(a) $2\sin(x + 30)$ incorrect equation		
(b) Incorrect translation: Sketch of $2\sin(x + 30)$			(b) Sketch of $2\sin(x + 30)$		
Only • ⁶ is available			All 3 marks are available		

Question			Generic scheme	Illustrative scheme	Max mark
15.	(a)		\bullet^1 state value of a \bullet^2 state value of b	\bullet^1 -5 \bullet^2 3	2
Notes:					
Commonly Observed Responses:					

Question			Generic scheme	Illustrative Scheme	Max Mark
15.	(b)		\bullet^3 state value of integral	\bullet^3 10	1
Notes:					
1. Candidates answer at (b) must be consistent with the value of b obtained in (a). 2. In parts (b) and (c), candidates who have 10 and -6 accompanied by working, the working must be checked to ensure that no errors have occurred prior to the correct answer appearing.					
Commonly Observed Responses:					
Candidate A From (a) $a = -3$ \bullet^1 ✗ $b = 5$ \bullet^2 ✗ $\int h(x)dx = 14$ \bullet^3 ✓1					

Question			Generic scheme	Illustrative scheme	Max mark
15.	(c)		\bullet^4 state value of derivative	\bullet^4 -6	1
Notes:					
Commonly Observed Responses:					

[END OF MARKING INSTRUCTIONS]



National
Qualifications
2017

2017 Mathematics Paper 2

Higher

Finalised Marking Instructions

© Scottish Qualifications Authority 2017

The information in this publication may be reproduced to support SQA qualifications only on a non-commercial basis. If it is reproduced, SQA should be clearly acknowledged as the source. If it is to be used for any other purpose, written permission must be obtained from permissions@sqa.org.uk.

Where the publication includes materials from sources other than SQA (secondary copyright), this material should only be reproduced for the purposes of examination or assessment. If it needs to be reproduced for any other purpose it is the centre's responsibility to obtain the necessary copyright clearance. SQA's NQ Assessment team may be able to direct you to the secondary sources.

These marking instructions have been prepared by examination teams for use by SQA appointed markers when marking external course assessments. This publication must not be reproduced for commercial or trade purposes.

General marking principles for Higher Mathematics

This information is provided to help you understand the general principles you must apply when marking candidate responses to questions in this Paper. These principles must be read in conjunction with the detailed marking instructions, which identify the key features required in candidate responses.

For each question the marking instructions are generally in two sections, namely Illustrative Scheme and Generic Scheme. The illustrative scheme covers methods which are commonly seen throughout the marking. The generic scheme indicates the rationale for which each mark is awarded. In general, markers should use the illustrative scheme and only use the generic scheme where a candidate has used a method not covered in the illustrative scheme.

- (a) Marks for each candidate response must always be assigned in line with these general marking principles and the detailed marking instructions for this assessment.
- (b) Marking should always be positive. This means that, for each candidate response, marks are accumulated for the demonstration of relevant skills, knowledge and understanding: they are not deducted from a maximum on the basis of errors or omissions.
- (c) If a specific candidate response does not seem to be covered by either the principles or detailed marking instructions, and you are uncertain how to assess it, you must seek guidance from your Team Leader.
- (d) Credit must be assigned in accordance with the specific assessment guidelines.
- (e) One mark is available for each •. There are no half marks.
- (f) Working subsequent to an error must be **followed through**, with possible credit for the subsequent working, provided that the level of difficulty involved is approximately similar. Where, subsequent to an error, the working for a follow through mark has been eased, the follow through mark cannot be awarded.
- (g) As indicated on the front of the question paper, full credit should only be given where the solution contains appropriate working. Unless specifically mentioned in the marking instructions, a correct answer with no working receives no credit.
- (h) Candidates may use any mathematically correct method to answer questions except in cases where a particular method is specified or excluded.
- (i) As a consequence of an error perceived to be trivial, casual or insignificant, eg $6 \times 6 = 12$ candidates lose the opportunity of gaining a mark. However, note the second example in comment (j).
- (j)

- (j) Where a transcription error (paper to script or within script) occurs, the candidate should normally lose the opportunity to be awarded the next process mark, eg

This is a transcription error and so the mark is not awarded.		$x^2 + 5x + 7 = 9x + 4$
Eased as no longer a solution of a quadratic equation so mark is not awarded.	→	$x - 4x + 3 = 0$
Exceptionally this error is not treated as a transcription error as the candidate deals with the intended quadratic equation. The candidate has been given the benefit of the doubt and all marks awarded.	→	$x = 1$
	→	$x^2 + 5x + 7 = 9x + 4$
	→	$x - 4x + 3 = 0$
	→	$(x - 3)(x - 1) = 0$
	→	$x = 1 \text{ or } 3$

- (k) **Horizontal/vertical marking**

Where a question results in two pairs of solutions, this technique should be applied, but only if indicated in the detailed marking instructions for the question.

Example:

	• ⁵		• ⁶
• ⁵	$x = 2$	$x = -4$	
• ⁶	$y = 5$	$y = -7$	

Horizontal: • ⁵ $x = 2$ and $x = -4$	Vertical: • ⁵ $x = 2$ and $y = 5$
• ⁶ $y = 5$ and $y = -7$	• ⁶ $x = -4$ and $y = -7$

Markers should choose whichever method benefits the candidate, but **not** a combination of both.

- (l) In final answers, unless specifically mentioned in the detailed marking instructions, numerical values should be simplified as far as possible, eg:

$\frac{15}{12}$ must be simplified to $\frac{5}{4}$ or $1\frac{1}{4}$	$\frac{43}{1}$ must be simplified to 43
$\frac{15}{0.3}$ must be simplified to 50	$\frac{4}{\cancel{5}}/3$ must be simplified to $\frac{4}{15}$
$\sqrt{64}$ must be simplified to 8*	

*The square root of perfect squares up to and including 100 must be known.

- (m) Commonly Observed Responses (COR) are shown in the marking instructions to help mark common and/or non-routine solutions. CORs may also be used as a guide when marking similar non-routine candidate responses.

(n) Unless specifically mentioned in the marking instructions, the following should not be penalised:

- Working subsequent to a correct answer
- Correct working in the wrong part of a question
- Legitimate variations in numerical answers/algebraic expressions, eg angles in degrees rounded to nearest degree
- Omission of units
- Bad form (bad form only becomes bad form if subsequent working is correct), eg $(x^3 + 2x^2 + 3x + 2)(2x + 1)$ written as $(x^3 + 2x^2 + 3x + 2) \times 2x + 1$

$2x^4 + 4x^3 + 6x^2 + 4x + x^3 + 2x^2 + 3x + 2$ written as $2x^4 + 5x^3 + 8x^2 + 7x + 2$ gains full credit

- Repeated error within a question, but not between questions or papers
- (o) In any 'Show that...' question, where the candidate has to arrive at a required result, the last mark of that part is not available as a follow-through from a previous error unless specified in the detailed marking instructions.
- (p) All working should be carefully checked, even where a fundamental misunderstanding is apparent early in the candidate's response. Marks may still be available later in the question so reference must be made continually to the marking instructions. The appearance of the correct answer does not necessarily indicate that the candidate has gained all the available marks.
- (q) Scored-out working which has not been replaced should be marked where still legible. However, if the scored out working has been replaced, only the work which has not been scored out should be marked.
- (r) Where a candidate has made multiple attempts using the same strategy and not identified their final answer, mark all attempts and award the lowest mark.

Where a candidate has tried different valid strategies, apply the above ruling to attempts within each strategy and then award the highest resultant mark.

For example:

Strategy 1 attempt 1 is worth 3 marks.	Strategy 2 attempt 1 is worth 1 mark.
Strategy 1 attempt 2 is worth 4 marks.	Strategy 2 attempt 2 is worth 5 marks.
From the attempts using strategy 1, the resultant mark would be 3.	From the attempts using strategy 2, the resultant mark would be 1.

In this case, award 3 marks.

Question			Generic scheme	Illustrative scheme	Max mark
1.	(a)		<ul style="list-style-type: none"> •¹ find mid-point of BC •² calculate gradient of BC •³ use property of perpendicular lines •⁴ determine equation of line in a simplified form 	<ul style="list-style-type: none"> •¹ $(6, -1)$ •² $-\frac{2}{6}$ •³ 3 •⁴ $y = 3x - 19$ 	4

Notes:

1. •⁴ is only available as a consequence of using a perpendicular gradient **and** a midpoint.
2. The gradient of the perpendicular bisector must appear in simplified form at •³ or •⁴ stage for •³ to be awarded.
3. At •⁴, accept $3x - y - 19 = 0$, $3x - y = 19$ or any other rearrangement of the equation where the constant terms have been simplified.

Commonly Observed Responses:

Question		Generic scheme	Illustrative scheme	Max mark
1.	(b)	<ul style="list-style-type: none"> •⁵ use $m = \tan \theta$ •⁶ determine equation of AB 	<ul style="list-style-type: none"> •⁵ 1 •⁶ $y = x - 3$ 	2

Notes:

4. At •⁶, accept $y - x + 3 = 0$, $y - x = -3$ or any other rearrangement of the equation where the constant terms have been simplified.

Commonly Observed Responses:

Question		Generic scheme	Illustrative scheme	Max mark
1.	(c)	<ul style="list-style-type: none"> •⁷ find x or y coordinate •⁸ find remaining coordinate 	<ul style="list-style-type: none"> •⁷ $x = 8$ or $y = 5$ •⁸ $y = 5$ or $x = 8$ 	2

Notes:

Commonly Observed Responses:

Question			Generic scheme	Illustrative scheme	Max mark
2.	(a)		<p>Method 1</p> <ul style="list-style-type: none"> •¹ know to use $x=1$ in synthetic division •² complete division, interpret result and state conclusion 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $\begin{array}{r rrrr} 1 & 2 & -5 & 1 & 2 \\ & & 2 & & \end{array}$ •² $\begin{array}{r rrrr} 1 & 2 & -5 & 1 & 2 \\ & & 2 & -3 & -2 \\ \hline & 2 & -3 & -2 & 0 \end{array}$ Remainder = 0 $\therefore (x-1)$ is a factor 	2
			<p>Method 2</p> <ul style="list-style-type: none"> •¹ know to substitute $x=1$ •² complete evaluation, interpret result and state conclusion 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ $2(1)^3 - 5(1)^2 + (1) + 2$ •² $= 0 \therefore (x-1)$ is a factor 	2
			<p>Method 3</p> <ul style="list-style-type: none"> •¹ start long division and find leading term in quotient •² complete division, interpret result and state conclusion 	<p>Method 3</p> <ul style="list-style-type: none"> •¹ $\begin{array}{r} 2x^2 \\ (x-1) \overline{) 2x^3 - 5x^2 + x + 2} \end{array}$ •² $\begin{array}{r} 2x^2 - 3x - 2 \\ (x-1) \overline{) 2x^3 - 5x^2 + x + 2} \\ \underline{2x^3 - 2x^2} \\ -3x^2 + x \\ \underline{-3x^2 + 3x} \\ -2x + 2 \\ \underline{-2x + 2} \\ 0 \end{array}$ remainder = 0 $\therefore (x-1)$ is a factor 	2

Question	Generic scheme	Illustrative scheme	Max mark
Notes:			
1. Communication at \bullet^2 must be consistent with working at that stage i.e. a candidate's working must arrive legitimately at 0 before \bullet^2 can be awarded. 2. Accept any of the following for \bullet^2 : <ul style="list-style-type: none"> '$f(1) = 0$ so $(x-1)$ is a factor' 'since remainder = 0, it is a factor' the 0 from any method linked to the word 'factor' by e.g. 'so', 'hence', '\therefore', '\rightarrow', '\Rightarrow' 3. Do not accept any of the following for \bullet^2 : <ul style="list-style-type: none"> double underlining the zero or boxing the zero without comment '$x = -1$ is a factor', '$(x+1)$ is a factor', '$(x+1)$ is a root', '$x = 1$ is a root', '$(x-1)$ is a root' '$x = -1$ is a root'. the word 'factor' only with no link 			
Commonly Observed Responses:			

Question	Generic scheme	Illustrative scheme	Max mark
2. (b)	\bullet^3 state quadratic factor \bullet^4 find remaining factors \bullet^5 state solution	\bullet^3 $2x^2 - 3x - 2$ \bullet^4 $(2x+1)$ and $(x-2)$ \bullet^5 $x = -\frac{1}{2}, 1, 2$	3
Notes:			
4. The appearance of " $= 0$ " is not required for \bullet^5 to be awarded. 5. Candidates who identify a different initial factor and subsequent quadratic factor can gain all available marks. 6. \bullet^5 is only available as a result of a valid strategy at \bullet^3 and \bullet^4 . 7. Accept $\left(-\frac{1}{2}, 0\right), (1, 0), (2, 0)$ for \bullet^5 .			
Commonly Observed Responses:			

Question			Generic scheme	Illustrative scheme	Max mark
3.			<ul style="list-style-type: none"> •¹ substitute for y •² express in standard quadratic form •³ factorise •⁴ find x coordinates •⁵ find y coordinates 	<ul style="list-style-type: none"> •¹ $(x-2)^2 + (3x-1)^2 = 25$ or $x^2 - 4x + 4 + (3x)^2 - 2(3x) + 1 = 25$ •² $10x^2 - 10x - 20 = 0$ •³ $10(x-2)(x+1) = 0$ <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <ul style="list-style-type: none"> •⁴ $x = 2$ •⁵ $y = 6$ </div> <div style="text-align: center;"> <ul style="list-style-type: none"> •⁵ $x = -1$ $y = -3$ </div> </div> 	5

Notes:

1. At •³ the quadratic must lead to two distinct real roots for •⁴ and •⁵ to be available.
2. •² is only available if ' = 0 ' appears at •² or •³ stage.
3. If a candidate arrives at an equation which is not a quadratic at •² stage, then •³, •⁴ and •⁵ are not available
4. At •³ do not penalise candidates who fail to extract the common factor or who have divided the quadratic equation by 10.
5. •³ is available for substituting correctly into the quadratic formula.
6. •⁴ and •⁵ may be marked either horizontally or vertically.
7. For candidates who identify **both** solutions by inspection, full marks may be awarded provided they justify that their points lie on **both** the line and the circle. Candidates who identify **both** solutions, but justify only one gain 2 out of 5.

Commonly Observed Responses:

Candidate A

$$(x-2)^2 + (3x-1)^2 = 25 \quad \bullet^1 \checkmark$$

$$10x^2 - 10x = 20 \quad \bullet^2 \times$$

$$10x(x-1) = 20 \quad \bullet^3 \boxed{\checkmark 2}$$

$$x = 2 \quad x = 3 \quad \bullet^4 \times$$

$$y = 6 \quad y = 9 \quad \bullet^5 \boxed{\checkmark 2}$$

Candidate B

Candidates who substitute into the circle equation only

$$\bullet^1 \checkmark$$

$$\bullet^2 \checkmark$$

$$\bullet^3 \checkmark$$

$$\bullet^4 \checkmark$$

Sub $x = 2$

$$y^2 - 2y - 24 = 0$$

$$(y-6)(y+4) = 0$$

$$y = 6 \text{ or } y = -4$$

Sub $x = -1$

$$y^2 - 2y - 15 = 0$$

$$(y+3)(y-5) = 0$$

$$y = -3 \text{ or } y = 5$$

(2, 6) (-1, -3) $\bullet^5 \times$

Question			Generic scheme	Illustrative scheme	Max mark
4.	(a)		Method 1 <ul style="list-style-type: none"> •¹ identify common factor •² complete the square •³ process for c and write in required form 	Method 1 <ul style="list-style-type: none"> •¹ $3(x^2 + 8x \dots \dots)$ stated or implied by •² •² $3(x+4)^2 \dots \dots$ •³ $3(x+4)^2 + 2$ 	3
			Method 2 <ul style="list-style-type: none"> •¹ expand completed square form •² equate coefficients •³ process for b and c and write in required form 	Method 2 <ul style="list-style-type: none"> •¹ $ax^2 + 2abx + ab^2 + c$ •² $a = 3, 2ab = 24, ab^2 + c = 50$ •³ $3(x+4)^2 + 2$ 	3

Notes:

1. $3(x+4)^2 + 2$ with no working gains •¹ and •² only; however, see Candidate G.
2. •³ is only available for a calculation involving both multiplication and subtraction of integers.

Commonly Observed Responses:

Candidate A $3\left(x^2 + 8x + \frac{50}{3}\right)$ $3\left(x^2 + 8x + 16 - 16 + \frac{50}{3}\right)$ <ul style="list-style-type: none"> •¹ ✓ •² ^ further working is required 	Candidate B $3x^2 + 24x + 50 = 3(x+8)^2 - 64 + 50$ $= 3(x+8)^2 - 14$ <ul style="list-style-type: none"> •¹ ✗ •² ✗ •³ ✓2
Candidate C $ax^2 + 2abx + ab^2 + c$ $a = 3, 2ab = 24, b^2 + c = 50$ $a = 3, b = 4, c = 34$ $3(x+4)^2 + 34$ <ul style="list-style-type: none"> •¹ ✓ •² ✗ •³ ✓1 	Candidate D $3((x^2 + 24x) + 50)$ $3((x+12)^2 - 144) + 50$ $3(x+12)^2 - 382$ <ul style="list-style-type: none"> •¹ ✗ •² ✓1 •³ ✓1

Question	Generic scheme	Illustrative scheme	Max mark
Candidate E $a(x+b)^2 + c = ax^2 + 2abx + ab^2 + c$ • ¹ ✓ $a = 3, 2ab = 24, ab^2 + c = 50$ • ² ✓ $b = 4, c = 2$ • ³ ✓ <div>•³ is awarded as all working relates to completed square form</div>	Candidate F $ax^2 + 2abx + ab^2 + c$ • ¹ ✓ $a = 3, 2ab = 24, ab^2 + c = 50$ • ² ✓ $b = 4, c = 2$ • ³ ✗ <div>•³ is lost as no reference is made to completed square form</div>		
Candidate G $3(x+4)^2 + 2$ Check: $3(x^2 + 8x + 16) + 2$ $= 3x^2 + 24x + 48 + 2$ $= 3x^2 + 24x + 50$ Award 3/3	Candidate H $3x^2 + 24x + 50$ $= 3(x+4)^2 - 16 + 50$ • ¹ ✓ • ² ✓ $= 3(x+4)^2 + 34$ • ³ ✗		

Question	Generic scheme	Illustrative scheme	Max mark
4. (b)	• ⁴ differentiate two terms • ⁵ complete differentiation	• ⁴ $3x^2 + 24x....$ • ⁵ $.... + 50$	2
Notes:			
3. • ⁴ is awarded for any two of the following three terms: $3x^2$, $+24x$, $+50$			
Commonly Observed Responses:			

Question			Generic scheme	Illustrative scheme	Max mark
4.	(c)		<p>Method 1</p> <ul style="list-style-type: none"> •⁶ link with (a) and identify sign of $(x+4)^2$ •⁷ communicate reason <p>Method 2</p> <ul style="list-style-type: none"> •⁶ identify minimum value of $f'(x)$ •⁷ communicate reason 	<p>Method 1</p> <ul style="list-style-type: none"> •⁶ $f'(x) = 3(x+4)^2 + 2$ and $(x+4)^2 \geq 0 \forall x$ •⁷ $\therefore 3(x+4)^2 + 2 > 0 \Rightarrow$ always strictly increasing <p>Method 2</p> <ul style="list-style-type: none"> •⁶ eg minimum value = 2 or annotated sketch •⁷ $2 > 0 \therefore (f'(x) > 0) \Rightarrow$ always strictly increasing 	2

Notes:

- Do not penalise $(x+4)^2 > 0$ or the omission of $f'(x)$ at •⁶ in Method 1.
- Responses in part (c) must be consistent with working in parts (a) and (b) for •⁶ and •⁷ to be available.
- Where erroneous working leads to a candidate considering a function which is not always strictly increasing, only •⁶ is available.
- At •⁶ communication should be explicitly in terms of the given function. Do not accept statements such as “(something)² ≥ 0 ”, “something squared ≥ 0 ”. However, •⁷ is still available.

Commonly Observed Responses:

<p>Candidate I</p> $f'(x) = 3(x+4)^2 + 2$ $3(x+4)^2 + 2 > 0 \Rightarrow \text{strictly increasing.}$ <p>Award 1 out of 2</p>	<p>Candidate J</p> <p>Since $3x^2 + 24x + 50 = 3(x+4)^2 + \frac{166}{50}$</p> <p>and $(x+4)^2$ is > 0 for all x then</p> $3(x+4)^2 + \frac{166}{50} > 0 \text{ for all } x.$ <p>Hence the curve is strictly increasing for all values of x. •⁶ ✓ •⁷ ✓1</p>
---	---

Question			Generic scheme	Illustrative scheme	Max mark
5.	(a)		• ¹ identify pathway • ² state \overrightarrow{PQ}	• ¹ $\overrightarrow{PR} + \overrightarrow{RQ}$ stated or implied by • ² • ² $-3\mathbf{i} - 4\mathbf{j} + 5\mathbf{k}$	2

Notes:

- Award •¹ $(9\mathbf{i} + 5\mathbf{j} + 2\mathbf{k}) + (-12\mathbf{i} - 9\mathbf{j} + 3\mathbf{k})$.
- Candidates who choose to work with column vectors and leave their answer in the form $\begin{pmatrix} -3 \\ -4 \\ 5 \end{pmatrix}$ cannot gain •².
- ² is not available for simply adding or subtracting vectors within an invalid strategy.
- Where candidates choose specific points consistent with the given vectors, only •¹ and •⁴ are available. However, should the statement 'without loss of generality' precede the selected points then marks •¹, •², •³ and •⁴ are all available.

Commonly Observed Responses:

Question			Generic scheme	Illustrative scheme	Max mark
5.	(b)		• ³ interpret ratio • ⁴ identify pathway and demonstrate result	• ³ $\frac{2}{3}$ or $\frac{1}{3}$ • ⁴ $\overrightarrow{PR} + \frac{2}{3}\overrightarrow{RQ}$ or $\overrightarrow{PQ} + \frac{1}{3}\overrightarrow{QR}$ leading to $\mathbf{i} - \mathbf{j} + 4\mathbf{k}$	2

Notes:

- This is a 'show that' question. Candidates who choose to work with column vectors must write their final answer in the required form to gain •⁴. $\begin{pmatrix} 1 \\ -1 \\ 4 \end{pmatrix}$ does not gain •⁴.
- Beware of candidates who fudge their working between •³ and •⁴.

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
Candidate A - legitimate use of the section formula		Candidate B - BEWARE - treating P as the origin	
$\vec{PS} = \frac{n\vec{PQ} + m\vec{PR}}{m+n}$ $\vec{PS} = \frac{2\vec{PQ} + \vec{PR}}{3} \quad \bullet^3 \checkmark$ $\vec{PS} = \frac{2 \begin{pmatrix} -3 \\ -4 \\ 5 \end{pmatrix} + \begin{pmatrix} 9 \\ 5 \\ 2 \end{pmatrix}}{3} + \frac{\begin{pmatrix} 9 \\ 5 \\ 2 \end{pmatrix}}{3}$ $= \begin{pmatrix} -2 \\ -8/3 \\ 10/3 \end{pmatrix} + \begin{pmatrix} 3 \\ 5/3 \\ 2/3 \end{pmatrix}$ $= \begin{pmatrix} 1 \\ -1 \\ 4 \end{pmatrix}$ $\vec{PS} = \mathbf{i} - \mathbf{j} + 4\mathbf{k} \quad \bullet^4 \checkmark$		$2\vec{QS} = \vec{SR}$ $3\mathbf{s} = 2\mathbf{q} + \mathbf{r} \quad \bullet^3 \checkmark$ $3\mathbf{s} = 2 \begin{pmatrix} -3 \\ -4 \\ 5 \end{pmatrix} + \begin{pmatrix} 9 \\ 5 \\ 2 \end{pmatrix}$ $\mathbf{s} = \mathbf{i} - \mathbf{j} + 4\mathbf{k} \quad \bullet^4 \times$	

Question			Generic scheme	Illustrative scheme	Max mark
5.	(c)		<p>Method 1</p> <ul style="list-style-type: none"> •⁵ evaluate $\overrightarrow{PQ} \cdot \overrightarrow{PS}$ •⁶ evaluate \overrightarrow{PQ} •⁷ evaluate \overrightarrow{PS} •⁸ use scalar product •⁹ calculate angle 	<p>Method 1</p> <ul style="list-style-type: none"> •⁵ $\overrightarrow{PQ} \cdot \overrightarrow{PS} = 21$ •⁶ $\overrightarrow{PQ} = \sqrt{50}$ •⁷ $\overrightarrow{PS} = \sqrt{18}$ •⁸ $\cos QPS = \frac{21}{\sqrt{50} \times \sqrt{18}}$ •⁹ 45.6° or 0.795 radians 	5
			<p>Method 2</p> <ul style="list-style-type: none"> •⁵ evaluate \overrightarrow{QS} •⁶ evaluate \overrightarrow{PQ} •⁷ evaluate \overrightarrow{PS} •⁸ use cosine rule •⁹ calculate angle 	<p>Method 2</p> <ul style="list-style-type: none"> •⁵ $\overrightarrow{QS} = \sqrt{26}$ •⁶ $\overrightarrow{PQ} = \sqrt{50}$ •⁷ $\overrightarrow{PS} = \sqrt{18}$ •⁸ $\cos QPS = \frac{(\sqrt{50})^2 + (\sqrt{18})^2 - (\sqrt{26})^2}{2 \times \sqrt{50} \times \sqrt{18}}$ •⁹ 45.6° or 0.795 radians 	5
Notes: <p>7. For candidates who use \overrightarrow{PS} not equal to $\mathbf{i} - \mathbf{j} + 4\mathbf{k}$ •⁵ is not available in Method 1 or •⁷ in Method 2.</p> <p>8. Do not penalise candidates who treat negative signs with a lack of rigour when calculating a magnitude. However, $\sqrt{1^2 - 1^2 + 4^2}$ leading to $\sqrt{16}$ indicates an invalid method for calculating the magnitude. No mark can be awarded for any magnitude arrived at using an invalid method.</p> <p>9. •⁸ is not available to candidates who simply state the formula $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{ \mathbf{a} \mathbf{b} }$.</p> <p>However, $\cos \theta = \frac{\overrightarrow{PQ} \cdot \overrightarrow{PS}}{ \overrightarrow{PQ} \times \overrightarrow{PS} }$ or $\cos \theta = \frac{21}{\sqrt{50} \times \sqrt{18}}$ is acceptable. Similarly for Method 2.</p> <p>10. Accept answers which round to 46° or 0.8 radians.</p> <p>11. Do not penalise the omission or incorrect use of units.</p> <p>12. •⁹ is only available as a result of using a valid strategy.</p> <p>13. •⁹ is only available for a single angle.</p> <p>14. For a correct answer with no working award 0/5.</p>					

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
Candidate C - Calculating wrong angle		Candidate D- Calculating wrong angle	
$\overrightarrow{QP} \cdot \overrightarrow{QS} = 29$ • ⁵ ✗ $ \overrightarrow{QP} = \sqrt{50}$ • ⁶ ✓1 $ \overrightarrow{QS} = \sqrt{26}$ • ⁷ ✓1 $\cos \hat{PQS} = \frac{29}{\sqrt{50} \times \sqrt{26}}$ • ⁸ ✓1 $\hat{PQS} = 36.5$ • ⁹ ✗ strategy incomplete		$\overrightarrow{PS} \cdot \overrightarrow{QP} = -21$ • ⁵ ✗ $ \overrightarrow{QP} = \sqrt{50}$ • ⁶ ✓ $ \overrightarrow{PS} = \sqrt{18}$ • ⁷ ✓ $\cos \theta = \frac{-21}{\sqrt{50} \times \sqrt{18}}$ • ⁸ ✓1 $\theta = 134.4$ • ⁹ ✗ strategy incomplete	
For candidates who continue, and use the angle found to evaluate the required angle, then all marks are available.		For candidates who continue, and use the angle found to evaluate the required angle, then all marks are available.	
Candidate E From (a) $\overrightarrow{PQ} = -21\mathbf{i} - 14\mathbf{j} + \mathbf{k}$		Candidate F From (a) $\overrightarrow{PQ} = 21\mathbf{i} + 14\mathbf{j} - \mathbf{k}$	
$\overrightarrow{PQ} \cdot \overrightarrow{PS} = -3$ • ⁵ ✓1 $ \overrightarrow{PQ} = \sqrt{638}$ • ⁶ ✓1 $ \overrightarrow{PS} = \sqrt{18}$ • ⁷ ✓ $\cos \hat{QPS} = \frac{-3}{\sqrt{638} \times \sqrt{18}}$ • ⁸ ✓1 $\hat{QPS} = 91.6$ • ⁹ ✓1		$\overrightarrow{PQ} \cdot \overrightarrow{PS} = 3$ • ⁵ ✓1 $ \overrightarrow{PQ} = \sqrt{638}$ • ⁶ ✓1 $ \overrightarrow{PS} = \sqrt{18}$ • ⁷ ✓ $\cos \hat{QPS} = \frac{3}{\sqrt{638} \times \sqrt{18}}$ • ⁸ ✓1 $\hat{QPS} = 88.4$ • ⁹ ✓1	
Candidate G From (b) $\overrightarrow{PS} = -4\mathbf{i} - 3\mathbf{j} + \mathbf{k}$			
$\overrightarrow{PQ} \cdot \overrightarrow{PS} = 3$ • ⁵ ✗ $ \overrightarrow{PQ} = \sqrt{50}$ • ⁶ ✓ $ \overrightarrow{PS} = \sqrt{26}$ • ⁷ ✓1 $\cos \hat{QPS} = \frac{3}{\sqrt{50} \times \sqrt{26}}$ • ⁸ ✓1 $\hat{QPS} = 85.2$ • ⁹ ✓1			

Question			Generic scheme	Illustrative scheme	Max mark
6.			<ul style="list-style-type: none"> •¹ substitute appropriate double angle formula •² express in standard quadratic form •³ factorise •⁴ solve for $\sin x^\circ$ •⁵ solve for x 	<ul style="list-style-type: none"> •¹ $5\sin x - 4 = 2(1 - 2\sin^2 x)$ •² $4\sin^2 x + 5\sin x - 6 = 0$ •³ $(4\sin x - 3)(\sin x + 2)$ <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <ul style="list-style-type: none"> •⁴ •⁴ $\sin x = \frac{3}{4},$ </div> <div style="text-align: center;"> <ul style="list-style-type: none"> •⁵ $\sin x = -2$ </div> </div> •⁵ $x = 0.848, 2.29, \cancel{\sin x = -2}$ 	5

Notes:

1. •¹ is not available for simply stating $\cos 2x = 1 - 2\sin^2 x$ with no further working.
2. In the event of $\cos^2 x^\circ - \sin^2 x^\circ$ or $2\cos^2 x^\circ - 1$ being substituted for $\cos 2x$, •¹ cannot be awarded until the equation reduces to a quadratic in $\sin x^\circ$.
3. Substituting $1 - 2\sin^2 A$ or $1 - 2\sin^2 \alpha$ for $\cos 2x$ at •¹ stage should be treated as bad form provided the equation is written in terms of x at •² stage. Otherwise, •¹ is not available.
4. '= 0' must appear by •³ stage for •² to be awarded. However, for candidates using the quadratic formula to solve the equation, '= 0' must appear at •² stage for •² to be awarded.
5. $5\sin x + 4\sin^2 x - 6 = 0$ does not gain •² unless •³ is awarded.
6. $\sin x = \frac{-5 \pm \sqrt{121}}{8}$ gains •³.
7. Candidates may express the equation obtained at •² in the form $4s^2 + 5s - 6 = 0$ or $4x^2 + 5x - 6 = 0$. In these cases, award •³ for $(4s - 3)(s + 2) = 0$ or $(4x - 3)(x + 2) = 0$. However, •⁴ is only available if $\sin x$ appears explicitly at this stage.
8. •⁴ and •⁵ are only available as a consequence of solving a quadratic equation.
9. •³, •⁴ and •⁵ are not available for any attempt to solve a quadratic equation written in the form $ax^2 + bx = c$.
10. •⁵ is not available to candidates who work in degrees and do not convert their solutions into radian measure.
11. Accept answers which round to 0.85 and 2.3 at •⁵ eg $\frac{49\pi}{180}, \frac{131\pi}{180}$.
12. Answers written as decimals should be rounded to no fewer than 2 significant figures.
13. Do not penalise additional solutions at •⁵.

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
Candidate A		Candidate B	
$\bullet^1 \checkmark \bullet^2 \checkmark$ $(4s-3)(s+2) = 0$ $s = \frac{3}{4}, s = -2$ $x = 0.848, 2.29$	$\bullet^3 \checkmark$ $\bullet^4 \times$ $\bullet^5 \checkmark$	$\bullet^1 \checkmark$ $4\sin^2 x + 5\sin x - 6 = 0$ $9\sin x - 6 = 0$ $\sin x = \frac{2}{3}$ $x = 0.730, 2.41$	$\bullet^2 \checkmark$ $\bullet^3 \times$ $\bullet^4 \boxed{\checkmark 2}$ $\bullet^5 \boxed{\checkmark 2}$
Candidate C		Candidate D	
$5\sin x - 4 = 2(1 - 2\sin^2 x)$ $4\sin^2 x + 5\sin x = 6$ $\sin x(4\sin x + 5) = 6$ $\sin x = 6, 4\sin x + 5 = 6$ no solution, $\sin x = \frac{1}{4}$ $x = 0.253, 2.89$	$\bullet^1 \checkmark$ $\bullet^2 \boxed{\checkmark 2}$ $\bullet^3 \boxed{\checkmark 2}$ $\bullet^4 \times$ $\bullet^5 \times$	$5\sin x - 4 = 2(1 - 2\sin^2 x)$ $4\sin^2 x + 5\sin x - 6 = 0$ $4\sin^2 x + 5\sin x = 6$ $\sin x(4\sin x + 5) = 6$ $\sin x = 6, 4\sin x + 5 = 6$ no solution, $\sin x = \frac{1}{4}$ $x = 0.253, 2.89$	$\bullet^1 \checkmark$ $\bullet^2 \checkmark$ $\bullet^3 \boxed{\checkmark 2}$ $\bullet^4 \times$ $\bullet^5 \times$
Candidate E - reading $\cos 2x$ as $\cos^2 x$			
$5\sin x - 4 = 2\cos^2 x$ $5\sin x - 4 = 2(1 - \sin^2 x)$ $2\sin^2 x + 5\sin x - 6 = 0$ $\sin x = \frac{-5 \pm \sqrt{73}}{4}$ $\sin x = 0.886, \sin x = -3.386$ $x = 1.08, 2.05$	$\bullet^1 \times$ $\bullet^2 \boxed{\checkmark 1}$ $\bullet^3 \boxed{\checkmark 1}$ $\bullet^4 \boxed{\checkmark 1}$ $\bullet^5 \boxed{\checkmark 1}$		

Question			Generic scheme	Illustrative scheme	Max mark
7.	(a)		<ul style="list-style-type: none"> •¹ write in differentiable form •² differentiate one term •³ complete differentiation and equate to zero •⁴ solve for x 	<ul style="list-style-type: none"> •¹ $\dots - 2x^{\frac{3}{2}}$ stated or implied •² $\frac{dy}{dx} = 6\dots$ or $\frac{dy}{dx} = \dots - 3x^{\frac{1}{2}}\dots$ •³ $\dots - 3x^{\frac{1}{2}} = 0$ or $6\dots = 0$ •⁴ $x = 4$ 	4

Notes:

1. For candidates who do not differentiate a term involving a fractional index, either •² or •³ is available but not both.
2. •⁴ is available only as a consequence of solving an equation involving a fractional power of x .
3. For candidates who integrate one or other of the terms •⁴ is unavailable.

Commonly Observed Responses:

Candidate A - differentiating incorrectly

$$y = 6x - 2x^{\frac{3}{2}} \quad \bullet^1 \checkmark$$

$$\frac{dy}{dx} = 6 - 3x^{\frac{5}{2}} \quad \bullet^2 \checkmark$$

$$6 - 3x^{\frac{5}{2}} = 0 \quad \bullet^3 \times$$

$$x = 1.32 \quad \bullet^4 \boxed{\checkmark 1}$$

Candidate B - integrating the second term

$$y = 6x - 2x^{\frac{3}{2}} \quad \bullet^1 \checkmark$$

$$\frac{dy}{dx} = 6 - \frac{4}{5}x^{\frac{5}{2}} \quad \bullet^2 \checkmark$$

$$6 - \frac{4}{5}x^{\frac{5}{2}} = 0 \quad \bullet^3 \times$$

$$x = 2.24 \quad \bullet^4 \times$$

Question			Generic scheme	Illustrative scheme	Max mark
7.	(b)		<ul style="list-style-type: none"> •⁵ evaluate y at stationary point •⁶ consider value of y at end points •⁷ state greatest and least values 	<ul style="list-style-type: none"> •⁵ 8 •⁶ 4 and 0 •⁷ greatest 8, least 0 stated explicitly 	3

Notes:

4. The only valid approach to finding the stationary point is via differentiation. A numerical approach can only gain •⁶.
5. •⁷ is not available to candidates who do not consider both end points.
6. Vertical marking is not applicable to •⁶ and •⁷.
7. Ignore any nature table which may appear in a candidate's solution; however, the appearance of (4,8) at a nature table is sufficient for •⁵.
8. Greatest (4,8); least (9,0) does not gain •⁷.
9. •⁵ and •⁷ are not available for evaluating y at a value of x , obtained at •⁴ stage, which lies outwith the interval $1 \leq x \leq 9$.
10. For candidates who only evaluate the derivative, •⁵, •⁶ and •⁷ are not available.

Commonly Observed Responses:

Question			Generic scheme	Illustrative scheme	Max mark
8.	(a)		<ul style="list-style-type: none"> •¹ find expression for u_1 •² find expression for u_2 and express in the correct form 	<ul style="list-style-type: none"> •¹ $5k - 20$ •² $u_2 = k(5k - 20) - 20$ leading to $u_2 = 5k^2 - 20k - 20$ 	2

Notes:

Commonly Observed Responses:

Question			Generic scheme	Illustrative scheme	Max mark
8.	(b)		<ul style="list-style-type: none"> •³ interpret information •⁴ express inequality in standard quadratic form •⁵ determine zeros of quadratic expression •⁶ state range with justification 	<ul style="list-style-type: none"> •³ $5k^2 - 20k - 20 < 5$ •⁴ $5k^2 - 20k - 25 < 0$ •⁵ $-1, 5$ •⁶ $-1 < k < 5$ with eg sketch or table of signs 	4
Notes:					
<p>1. Candidates who work with an equation from the outset lose •³ and •⁴. However, •⁵ and •⁶ are still available.</p> <p>2. At •⁵ do not penalise candidates who fail to extract the common factor or who have divided the quadratic inequation by 5.</p> <p>3. •⁴ and •⁵ are only available to candidates who arrive at a quadratic expression at •³.</p> <p>4. At •⁶ accept “$k > -1$ and $k < 5$” or “$k > -1, k < 5$” together with the required justification.</p> <p>5. For a trial and error approach award 0/4.</p>					
Commonly Observed Responses:					

Question			Generic scheme	Illustrative scheme	Max mark
9.			<p>Method 1</p> <ul style="list-style-type: none"> •¹ state linear equation •² introduce logs •³ use laws of logs •⁴ use laws of logs •⁵ state k and n 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $\log_2 y = \frac{1}{4}\log_2 x + 3$ •² $\log_2 y = \frac{1}{4}\log_2 x + 3\log_2 2$ •³ $\log_2 y = \log_2 x^{\frac{1}{4}} + \log_2 2^3$ •⁴ $\log_2 y = \log_2 2^3 x^{\frac{1}{4}}$ •⁵ $k = 8, n = \frac{1}{4}$ or $y = 8x^{\frac{1}{4}}$ 	5
			<p>Method 2</p> <ul style="list-style-type: none"> •¹ state linear equation •² use laws of logs •³ use laws of logs •⁴ use laws of logs •⁵ state k and n 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ $\log_2 y = \frac{1}{4}\log_2 x + 3$ •² $\log_2 y = \log_2 x^{\frac{1}{4}} + 3$ •³ $\log_2 \frac{y}{x^{\frac{1}{4}}} = 3$ •⁴ $\frac{y}{x^{\frac{1}{4}}} = 2^3$ •⁵ $k = 8, n = \frac{1}{4}$ or $y = 8x^{\frac{1}{4}}$ 	5

Question			Generic Scheme	Illustrative Scheme	Max Mark
			<p>Method 3</p> <ul style="list-style-type: none"> •¹ introduce logs to $y = kx^n$ •² use laws of logs •³ interpret intercept •⁴ use laws of logs •⁵ interpret gradient 	<p>Method 3</p> <p>The equations at •¹, •² and •³ must be stated explicitly.</p> <ul style="list-style-type: none"> •¹ $\log_2 y = \log_2 kx^n$ •² $\log_2 y = n \log_2 x + \log_2 k$ •³ $\log_2 k = 3$ •⁴ $k = 8$ •⁵ $n = \frac{1}{4}$ 	5
			<p>Method 4</p> <ul style="list-style-type: none"> •¹ interpret point on log graph •² convert from log to exp. form •³ interpret point and convert •⁴ substitute into $y = kx^n$ and evaluate k •⁵ substitute other point into $y = kx^n$ and evaluate n 	<p>Method 4</p> <ul style="list-style-type: none"> •¹ $\log_2 x = -12$ and $\log_2 y = 0$ •² $x = 2^{-12}$ and $y = 2^0$ •³ $\log_2 x = 0, \log_2 y = 3$ $x = 1, y = 2^3$ •⁴ $2^3 = k \times 1^n \Rightarrow k = 8$ •⁵ $2^0 = 2^3 \times 2^{-12n}$ $\Rightarrow 3 - 12n = 0$ $\Rightarrow n = \frac{1}{4}$ 	5
Notes: <ol style="list-style-type: none"> 1. Markers must not pick and choose between methods. Identify the method which best matches the candidates approach. 2. Treat the omission of base 2 as bad form at •¹ and •³ in Method 1, at •¹ and •² for Method 2 and Method 3, and at •¹ in Method 4. 3. '$m = \frac{1}{4}$', or 'gradient = $\frac{1}{4}$', does not gain •⁵ in Method 3. 4. Accept 8 in lieu of 2^3 throughout. 5. In Method 4 candidates may use (0,3) for •¹ and •² followed by (-12,0) for •³. 					

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
Candidate A With no working. Method 3: $k = 8$ • ⁴ ✓ $n = \frac{1}{4}$ • ⁵ ✓ Award 2/5		Candidate B With no working. Method 3: $n = 8$ • ⁴ ✗ $k = \frac{1}{4}$ • ⁵ ✗ Award 0/5	
Candidate C Method 3: $\log_2 k = 3$ • ³ ✓ $k = 8$ • ⁴ ✓ $n = \frac{1}{4}$ • ⁵ ✓ Award 3/5		Candidate D Method 2: $\log_2 y = \frac{1}{4} \log_2 x + 3$ • ¹ ✓ $\log_2 y = \log_2 x^{\frac{1}{4}} + 3$ • ² ✓ $y = x^{\frac{1}{4}} + 3$ • ³ ✗ • ⁴ ✗ $k = 1, n = \frac{1}{4}$ • ⁵ ✗ Award 2/5	
Candidate E Method 2: $y = \frac{1}{4}x + 3$ $y = \frac{1}{4}x + 3$ $\log_2 y = \frac{1}{4} \log_2 x + 3$ • ¹ ✓ $\log_2 y = \log_2 x^{\frac{1}{4}} + 3$ • ² ✓ $\frac{y}{x^{\frac{1}{4}}} = 3$ • ³ ^ • ⁴ ✗ $y = 3x^{\frac{1}{4}}$ • ⁵ ✓1 Award 3/5			

Question			Generic scheme	Illustrative scheme	Max mark
10.	(a)		Method 1 <ul style="list-style-type: none"> •¹ calculate m_{AB} •² calculate m_{BC} •³ interpret result and state conclusion 	Method 1 <ul style="list-style-type: none"> •¹ $m_{AB} = \frac{3}{9} = \frac{1}{3}$ see Note 1 •² $m_{BC} = \frac{5}{15} = \frac{1}{3}$ •³ ... \Rightarrow AB and BC are parallel (common direction), B is a common point, hence A, B and C are collinear. 	3
			Method 2 <ul style="list-style-type: none"> •¹ calculate an appropriate vector e.g. \overrightarrow{AB} •² calculate a second vector e.g. \overrightarrow{BC} and compare •³ interpret result and state conclusion 	Method 2 <ul style="list-style-type: none"> •¹ $\overrightarrow{AB} = \begin{pmatrix} 9 \\ 3 \end{pmatrix}$ see Note 1 •² $\overrightarrow{BC} = \begin{pmatrix} 15 \\ 5 \end{pmatrix} \therefore \overrightarrow{AB} = \frac{3}{5}\overrightarrow{BC}$ •³ ... \Rightarrow AB and BC are parallel (common direction), B is a common point, hence A, B and C are collinear. 	3
			Method 3 <ul style="list-style-type: none"> •¹ calculate m_{AB} •² find equation of line and substitute point •³ communication 	Method 3 <ul style="list-style-type: none"> •¹ $m_{AB} = \frac{3}{9} = \frac{1}{3}$ •² eg, $y - 1 = \frac{1}{3}(x - 2)$ leading to $6 - 1 = \frac{1}{3}(17 - 2)$ •³ since C lies on line A, B and C are collinear 	
Notes: <ol style="list-style-type: none"> At •¹ and •² stage, candidates may calculate the gradients/vectors using any pair of points. •³ can only be awarded if a candidate has stated “parallel”, “common point” and “collinear”. Candidates who state “points A, B and C are parallel” or “m_{AB} and m_{BC} are parallel” do not gain •³. 					

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
<p>Candidate A</p> $m_{AB} = \frac{3}{9} = \frac{1}{3}$ $m_{BC} = \frac{5}{15}$ <p>\Rightarrow AB and BC are parallel , B is a common point, hence A, B and C are collinear.</p>	<p>Candidate B</p> $\begin{pmatrix} 9 \\ 3 \end{pmatrix}$ $\begin{pmatrix} 15 \\ 5 \end{pmatrix} \therefore \overrightarrow{AB} = \frac{5}{3} \overrightarrow{BC}$ <p>\Rightarrow AB and BC are parallel , B is a common point, hence A, B and C are collinear.</p>	<p>Candidate C</p> $\overrightarrow{AB} = \begin{pmatrix} 9 \\ 3 \end{pmatrix}$ $\overrightarrow{BC} = \begin{pmatrix} 15 \\ 5 \end{pmatrix} = 5 \begin{pmatrix} 3 \\ 1 \end{pmatrix} \text{ and}$ $\begin{pmatrix} 9 \\ 3 \end{pmatrix} = 3 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ <p>$\therefore \overrightarrow{AB} = \frac{5}{3} \overrightarrow{BC}$ ignore working subsequent to correct statement at \bullet^2. \Rightarrow AB and BC are parallel , B is a common point, hence A, B and C are collinear.</p>	<p>$\bullet^1 \checkmark$</p> <p>$\bullet^2 \wedge$</p> <p>$\bullet^3 \boxed{\checkmark 1}$</p> <p>$\bullet^1 \checkmark$</p> <p>$\bullet^2 \times$</p> <p>$\bullet^3 \boxed{\checkmark 1}$</p> <p>$\bullet^1 \checkmark$</p> <p>$\bullet^2 \checkmark$</p> <p>$\bullet^3 \checkmark$</p>

Question			Generic scheme	Illustrative scheme	Max mark
11.	(a)		Method 1 • ¹ substitute for $\sin 2x$ • ² simplify and factorise • ³ substitute for $1 - \cos^2 x$ and simplify	Method 1 • ¹ $\frac{2 \sin x \cos x}{2 \cos x} - \sin x \cos^2 x$ stated explicitly as above or in a simplified form of the above • ² $\sin x(1 - \cos^2 x)$ • ³ $\sin x \times \sin^2 x$ leading to $\sin^3 x$	3
			Method 2 • ¹ substitute for $\sin 2x$ • ² simplify and substitute for $\cos^2 x$ • ³ expand and simplify	Method 2 • ¹ $\frac{2 \sin x \cos x}{2 \cos x} - \sin x \cos^2 x$ stated explicitly as above or in a simplified form of the above • ² $\sin x - \sin x(1 - \sin^2 x)$ • ³ $\sin x - \sin x + \sin^3 x$ leading to $\sin^3 x$	3

Notes:

- ¹ is not available to candidates who simply quote $\sin 2x = 2 \sin x \cos x$ without substituting into the expression given on the LHS. See Candidate B
- In method 2 where candidates attempt •¹ and •² in the same line of working •¹ may still be awarded if there is an error at •².
- ³ is not available to candidates who work throughout with A in place of x .
- Treat multiple attempts which are not scored out as different strategies, and apply General Marking Principle (r).
- On the appearance of $\text{LHS} = 0$, the first available mark is lost; however, any further marks are still available.

Commonly Observed Responses:

Candidate A

$$\frac{2 \sin x \cos x}{2 \cos x} - \sin x \cos^2 x = \sin^3 x \quad \bullet^1 \checkmark$$

$$\sin x - \sin x \cos^2 x = \sin^3 x \quad \bullet^2 \wedge$$

$$1 - \cos^2 x = \sin^2 x \quad \bullet^3 \times$$

$$\sin^2 x = \sin^2 x$$

In proving the identity, candidates must work with both sides independently ie in each line of working the LHS must be equivalent to the line above.

Candidate B

$$\text{LHS} = \frac{\sin 2x}{2 \cos x} - \sin x \cos^2 x$$

$$\frac{\sin 2x}{2 \cos x} = \frac{2 \sin x \cos x}{2 \cos x} = \sin x$$

$$\sin x - \sin x \cos^2 x \quad \bullet^1 \checkmark$$

$$\sin x(1 - \cos^2 x) \quad \bullet^2 \checkmark$$

Question			Generic scheme	Illustrative scheme	Max mark
11.	(b)		<ul style="list-style-type: none"> •⁴ know to differentiate $\sin^3 x$ •⁵ start to differentiate •⁶ complete differentiation 	<ul style="list-style-type: none"> •⁴ $\frac{d}{dx}(\sin^3 x)$ •⁵ $3\sin^2 x \dots$ •⁶ $\dots \times \cos x$ 	3
Notes:					
Commonly Observed Responses:					

[END OF MARKING INSTRUCTIONS]