



National
Qualifications
2023

2023 Mathematics
Higher - Paper 2
Finalised Marking Instructions

© Scottish Qualifications Authority 2023

These marking instructions have been prepared by examination teams for use by SQA appointed markers when marking external course assessments.

The information in this document may be reproduced in support of SQA qualifications only on a non-commercial basis. If it is reproduced, SQA must be clearly acknowledged as the source. If it is to be reproduced for any other purpose, written permission must be obtained from permissions@sqa.org.uk.



General marking principles for Higher Mathematics

Always apply these general principles. Use them in conjunction with the detailed marking instructions, which identify the key features required in candidates' responses.

For each question, the marking instructions are generally in two sections:

- *generic scheme* – this indicates why each mark is awarded
- *illustrative scheme* – this covers methods which are commonly seen throughout the marking

In general, you should use the illustrative scheme. Only use the generic scheme where a candidate has used a method not covered in the illustrative scheme.

- Always use positive marking. This means candidates accumulate marks for the demonstration of relevant skills, knowledge and understanding; marks are not deducted for errors or omissions.
- If you are uncertain how to assess a specific candidate response because it is not covered by the general marking principles or the detailed marking instructions, you must seek guidance from your team leader.
- One mark is available for each •. There are no half marks.
- If a candidate's response contains an error, all working subsequent to this error must still be marked. Only award marks if the level of difficulty in their working is similar to the level of difficulty in the illustrative scheme.
- Only award full marks where the solution contains appropriate working. A correct answer with no working receives no mark, unless specifically mentioned in the marking instructions.
- Candidates may use any mathematically correct method to answer questions, except in cases where a particular method is specified or excluded.
- If an error is trivial, casual or insignificant, for example $6 \times 6 = 12$, candidates lose the opportunity to gain a mark, except for instances such as the second example in point (h) below.
- If a candidate makes a transcription error (question paper to script or within script), they lose the opportunity to gain the next process mark, for example

This is a transcription error and so the mark is not awarded.

This is no longer a solution of a quadratic equation, so the mark is not awarded.

$$x^2 + 5x + 7 = 9x + 4$$

$$x - 4x + 3 = 0$$

$$x = 1$$

The following example is an exception to the above

This error is not treated as a transcription error, as the candidate deals with the intended quadratic equation. The candidate has been given the benefit of the doubt and all marks awarded.

$$x^2 + 5x + 7 = 9x + 4$$

$$x - 4x + 3 = 0$$

$$(x - 3)(x - 1) = 0$$

$$x = 1 \text{ or } 3$$

(i) **Horizontal/vertical marking**

If a question results in two pairs of solutions, apply the following technique, but only if indicated in the detailed marking instructions for the question.

Example:

$$\begin{array}{cc} \bullet^5 & \bullet^6 \\ \bullet^5 & x = 2 \quad x = -4 \\ \bullet^6 & y = 5 \quad y = -7 \end{array}$$

Horizontal: $\bullet^5 x = 2$ and $x = -4$ Vertical: $\bullet^5 x = 2$ and $y = 5$
 $\bullet^6 y = 5$ and $y = -7$ $\bullet^6 x = -4$ and $y = -7$

You must choose whichever method benefits the candidate, **not** a combination of both.

(j) In final answers, candidates should simplify numerical values as far as possible unless specifically mentioned in the detailed marking instruction. For example

$$\begin{array}{ll} \frac{15}{12} \text{ must be simplified to } \frac{5}{4} \text{ or } 1\frac{1}{4} & \frac{43}{1} \text{ must be simplified to } 43 \\ \frac{15}{0.3} \text{ must be simplified to } 50 & \frac{4\cancel{5}}{3} \text{ must be simplified to } \frac{4}{15} \\ \sqrt{64} \text{ must be simplified to } 8^* & \end{array}$$

*The square root of perfect squares up to and including 144 must be known.

(k) Commonly Observed Responses (COR) are shown in the marking instructions to help mark common and/or non-routine solutions. CORs may also be used as a guide when marking similar non-routine candidate responses.

(l) Do not penalise candidates for any of the following, unless specifically mentioned in the detailed marking instructions:

- working subsequent to a correct answer
- correct working in the wrong part of a question
- legitimate variations in numerical answers/algebraic expressions, for example angles in degrees rounded to nearest degree
- omission of units
- bad form (bad form only becomes bad form if subsequent working is correct), for example
 $(x^3 + 2x^2 + 3x + 2)(2x + 1)$ written as
 $(x^3 + 2x^2 + 3x + 2) \times 2x + 1$
 $= 2x^4 + 5x^3 + 8x^2 + 7x + 2$ gains full credit
- repeated error within a question, but not between questions or papers

(m) In any 'Show that...' question, where candidates have to arrive at a required result, the last mark is not awarded as a follow-through from a previous error, unless specified in the detailed marking instructions.

- (n) You must check all working carefully, even where a fundamental misunderstanding is apparent early in a candidate's response. You may still be able to award marks later in the question so you must refer continually to the marking instructions. The appearance of the correct answer does not necessarily indicate that you can award all the available marks to a candidate.
- (o) You should mark legible scored-out working that has not been replaced. However, if the scored-out working has been replaced, you must only mark the replacement working.
- (p) If candidates make multiple attempts using the same strategy and do not identify their final answer, mark all attempts and award the lowest mark. If candidates try different valid strategies, apply the above rule to attempts within each strategy and then award the highest mark.

For example:

Strategy 1 attempt 1 is worth 3 marks.	Strategy 2 attempt 1 is worth 1 mark.
Strategy 1 attempt 2 is worth 4 marks.	Strategy 2 attempt 2 is worth 5 marks.
From the attempts using strategy 1, the resultant mark would be 3.	From the attempts using strategy 2, the resultant mark would be 1.

In this case, award 3 marks.

Question		Generic scheme	Illustrative scheme	Max mark
1.	(a)	<ul style="list-style-type: none"> •¹ find gradient of QR •² use property of perpendicular lines •³ determine equation of altitude 	<ul style="list-style-type: none"> •¹ $-\frac{1}{3}$ or $-\frac{5}{15}$ •² 3 •³ $y = 3x - 16$ 	3
Notes:				
<p>1. •³ is only available to candidates who find and use a perpendicular gradient.</p> <p>2. The gradient of the perpendicular bisector must appear in fully simplified form at •² or •³ stage for •³ to be awarded - see Candidate B.</p> <p>3. •³ is not available as a consequence of using the midpoint of QR and the point P.</p> <p>4. At •³, accept any arrangement of a candidate's equation where constant terms have been simplified.</p>				
Commonly Observed Responses:				
		Candidate A - BEWARE Correct equation from incorrect substitution $m = \frac{13 - (-2)}{3 - (8)} = 3$ $y = 3x - 16$ <ul style="list-style-type: none"> •¹ ✗ •² ^ •³ ✗ 	Candidate B - unsimplified gradient $m = -\frac{5}{15}$ $m_{\perp} = \frac{15}{5}$ $15x - 5y - 80 = 0$ <ul style="list-style-type: none"> •¹ ✓ •² ✓ •³ ^ 	
	(b)	<ul style="list-style-type: none"> •⁴ determine gradient of the line •⁵ use $m = \tan \theta$ to find the angle 	<ul style="list-style-type: none"> •⁴ $m = \frac{1}{2}$ or $\tan \theta = \frac{1}{2}$ •⁵ 26.6° or 0.4636... radians 	2
Notes:				
<p>5. Do not penalise the omission of units at •⁵.</p> <p>6. Accept any answers which round to 27° or 0.46 radians.</p> <p>7. For 27° or 0.46 radians without working award 2/2.</p> <p>8. Where candidates find the angle of the altitude or other sides with the positive direction of the x-axis only •⁵ is available.</p>				
Commonly Observed Responses:				
		Candidate C - no reference to tan $m = \frac{4}{8}$ 26.6° <ul style="list-style-type: none"> •⁴ ✓ •⁵ ✓ 	Candidate D - BEWARE $m = \frac{1}{2}$ $\theta = \tan^{-1} \frac{1}{2}$ $\theta = 26.6^{\circ}$ Stating tan rather than \tan^{-1} See general marking principle (g) <ul style="list-style-type: none"> •⁴ ✓ •⁵ ✗ 	
		Candidate E $\tan^{-1}(3) = 72^{\circ}$ <ul style="list-style-type: none"> •⁴ ✗ •⁵ ✓₁ 		

Question	Generic scheme	Illustrative scheme	Max mark
2.	<ul style="list-style-type: none"> •¹ calculate y-coordinate •² differentiate •³ calculate the gradient •⁴ find equation of line 	<ul style="list-style-type: none"> •¹ -1 •² $10x^4 - 3$ •³ 7 •⁴ $y = 7x - 8$ 	4


Notes:

1. Only •¹ is available to candidates who integrate.
2. •⁴ is only available where candidates attempt to find the gradient by substituting into their derivative.
3. The appearance of $10x^4 - 3$ gains •².
4. •³ is not available for $y = 7$. However, where 7 is then used as the gradient of the straight line, •³ may be awarded - see Candidates B, C & D.
5. •⁴ is not available as a consequence of using a perpendicular gradient.

Commonly Observed Responses:

<p>Candidate A</p> $\frac{dy}{dx} = 10x^4 - 3$ $y = 7$ $m = -3$ $y = -3x + 10$	<ul style="list-style-type: none"> •² ✓ •¹ ✗ •³ ✗ •⁴ ✓₂ 	<p>Candidate B - incorrect notation</p> $y = -1$ $y = 10x^4 - 3$ $y = 7$ $y + 1 = 7(x - 1)$ $y = 7x - 8$	<ul style="list-style-type: none"> •¹ ✓ - BoD •² ✓ •³ ✓ - BoD •⁴ ✓
<p>Candidate C - use of values in equation</p> $y = -1$ $\frac{dy}{dx} = 10x^4 - 3$ $\frac{dy}{dx} = 7$ $y = 7$ $y + 1 = 7(x - 1)$ $y = 7x - 8$	<ul style="list-style-type: none"> •¹ ✓ - BoD •² ✓ •³ ✓ •⁴ ✓ 	<p>Candidate D - incorrect notation</p> $y = -1$ $\frac{dy}{dx} = 10x^4 - 3$ $y = 7$	<ul style="list-style-type: none"> •¹ ✓ - BoD •² ✓ •³ ✗ <p style="text-align: center;">Evidence for •³ would need to appear in the equation of the line</p>
<p>Candidate E</p> $y = -1$ $\frac{dy}{dx} = 10x^4 - 3 = 0$ $10(1)^4 - 3 = 0$ $m = 7$ $y = 7x - 8$	<ul style="list-style-type: none"> •¹ ✓ •² ✓ •³ ✗ •⁴ ✓₁ 		

Question		Generic scheme	Illustrative scheme	Max mark
3.		<ul style="list-style-type: none"> •¹ start to integrate •² complete integration 	<ul style="list-style-type: none"> •¹ $7 \sin\left(4x + \frac{\pi}{3}\right) \dots$ •² $\dots \times \frac{1}{4} + c$ 	2
Notes:				
<p>1. Award •¹ for any appearance of $(+)7 \sin\left(4x + \frac{\pi}{3}\right)$ regardless of any constant multiplier.</p> <p>2. Candidates who work in degrees from the start cannot gain •¹, however •² is still available - see Candidate C.</p> <p>3. Where candidates use any other invalid approach, eg $7 \sin\left(4x + \frac{\pi}{3}\right)^2$, $\int\left(7 \cos 4x + \cos \frac{\pi}{3}\right) dx$ or $7 \sin 4x + \frac{\pi}{3}$ award 0/2. However, see Candidate E.</p> <p>4. Do not penalise the appearance of an integral sign and/or dx throughout.</p>				
Commonly Observed Responses:				
Candidate A - using addition formula $\int\left(7 \cos 4x \cos \frac{\pi}{3} - 7 \sin 4x \sin \frac{\pi}{3}\right) dx$ $= \frac{7}{4} \sin 4x \cos \frac{\pi}{3} + \frac{7}{4} \cos 4x \sin \frac{\pi}{3} \dots \quad \bullet^1 \checkmark$ $= \frac{7}{4} \sin 4x \left(\frac{1}{2}\right) + \frac{7}{4} \cos 4x \left(\frac{\sqrt{3}}{2}\right) + c \quad \bullet^2 \checkmark$		Candidate B $\frac{7}{4} \sin\left(4x + \frac{\pi}{3}\right)$ $= \frac{7}{4} \sin\left(4x + \frac{\pi}{3}\right) + c$ <ul style="list-style-type: none"> •¹ ✓ •² ✓ 		
Candidate C - working in degrees $\int 7 \cos(4x + 60) dx$ $= 7 \sin(4x + 60) \times \frac{1}{4} + c \quad \bullet^1 \times \bullet^2 \boxed{\checkmark_1}$		Candidate D - integrating over two lines $7 \sin\left(4x + \frac{\pi}{3}\right)$ $= \frac{7}{4} \sin\left(4x + \frac{\pi}{3}\right) + c$ <ul style="list-style-type: none"> •¹ ✓ •² ✗ 		
Candidate E - integrating in part $-\frac{7}{4} \sin\left(4x + \frac{\pi}{3}\right) + c \quad \bullet^1 \times \bullet^2 \boxed{\checkmark_1}$		Candidate F - insufficient evidence of integration $\frac{7}{4} \cos\left(4x + \frac{\pi}{3}\right) + c$ <ul style="list-style-type: none"> •¹ ✗ •² ✗ 		

Question	Generic scheme	Illustrative scheme	Max mark
4.	<ul style="list-style-type: none"> •¹ reflect in the y-axis •² apply appropriate vertical scaling 	<ul style="list-style-type: none"> •¹ cubic graph with max at $(-2, 0)$ and passing through $(1, 0)$ •²  	2

Notes:

1. Where candidates do not sketch a cubic function award 0/2.
2. For transformations of the form $f(-x)+k$ or $-f(x+k)$ award 0/2.
3. If the transformation has not been applied to all coordinates, award 0/2.

Question	Generic scheme	Illustrative scheme	Max mark	
4.	(continued)			
Commonly Observed Responses:				
Function	Transformation of (-1,0) and (2,0)	Transformation of (0,-2)	Shape	Award
Incorrect orientation	(-2,0) and (1, 0)	(0,-4)		0/2
$-2f(x)$	(-1,0) and (2,0)	(0,4)		1/2
$-2f(-x)$	(-2,0) and (1, 0)	(0,4)		1/2
$-2f(-2x)$	(-1,0) and ($\frac{1}{2}$,0)	(0,4)		0/2
$-2f\left(-\frac{x}{2}\right)$	(-4,0) and (2,0)	(0,4)		0/2
$2f(x)$	(-1,0) and (2,0)	(0,-4)		1/2
$2f(2x)$	($-\frac{1}{2}$,0) and (1,0)	(0,-4)		1/2
$2f\left(\frac{x}{2}\right)$	(-2,0) and (4,0)	(0,-4)		1/2
$2f\left(-\frac{x}{2}\right)$	(-4,0) and (2,0)	(0,-4)		1/2
$2f(x-1)$	(0,0) and (3,0)	(1,-4)		1/2
$f(-x)$	(-2,0) and (1,0)	(0,-2)		1/2
$\frac{1}{2}f(-x)$	(-2,0) and (1,0)	(0,-1)		1/2
$f(2x)$	($-\frac{1}{2}$,0) and (1,0)	(0,-2)		0/2
$f(-2x)$	(-1,0) and ($\frac{1}{2}$,0)	(0,-2)		0/2
$f\left(-\frac{x}{2}\right)$	(-4,0) and (2,0)	(0,-2)		0/2
$-f\left(\frac{x}{2}\right)$	(-2,0) and (4,0)	(0,2)		0/2
$-f\left(-\frac{x}{2}\right)$	(-4,0) and (2,0)	(0,2)		0/2

Question	Generic scheme	Illustrative scheme	Max mark
5.	<ul style="list-style-type: none"> •¹ start to differentiate •² complete differentiation •³ calculate rate of change 	<ul style="list-style-type: none"> •¹ $4(3-2x)^3 \dots$ •² $\dots \times (-2)$ •³ 1000 	3

Notes:

1. Correct answer with no working, award 0/3.
2. Accept $4u^3 \times (-2)$ where $u = 3 - 2x$ for •¹.
3. Where candidates evaluate $f(4)$, award 0/3, see Candidate B.
4. •³ is only available for evaluating expressions equivalent to $k(3-2x)^3$.

Commonly Observed Responses:

<p>Candidate A</p> <p>$f'(x) = 4(3-2x)^3 \times (-2)$ •¹ ✓ •² ✓</p> <p>$f'(x) = 8(3-2x)^3$</p> <p>$f'(4) = -1000$ •³ ✗</p>	<p>Candidate B - evaluating f(x)</p> <p>$f'(x) = (3-2x)^4$ •¹ ✗ •² ✗</p> <p>$f'(4) = 625$ •³ ✗</p>
<p>Candidate C - differentiating over two lines</p> <p>$4(3-2x)^3$ •¹ ✓</p> <p>$4(3-2x)^3 \times 2$ •² ✗</p> <p>-1000 •³ <input checked="" type="checkbox"/> 1</p>	<p>Candidate D - differentiating over two lines</p> <p>$4(3-2x)^3$ •¹ ✓</p> <p>$4(3-2x)^3 \times -2$ •² ^</p> <p>1000 •³ <input checked="" type="checkbox"/> 1</p>
<p>Candidate E - insufficient evidence for mark 1</p> <p>$f'(x) = 8(3-2x)^3$ •¹ ✗ •² ✗</p> <p>$f'(4) = -1000$ •³ <input checked="" type="checkbox"/> 1</p>	<p>Candidate F</p> <p>$4(3-2x)^3$ •¹ ✓ •² ^</p> <p>-500 •³ <input checked="" type="checkbox"/> 1</p>

Question		Generic scheme	Illustrative scheme	Max mark
6.		<p>Method 1</p> <ul style="list-style-type: none"> •¹ equate composite function to x •² write $f(f^{-1}(x))$ in terms of $f^{-1}(x)$ •³ state inverse function 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $f(f^{-1}(x)) = x$ •² $x = \frac{2}{f^{-1}(x)} + 3$ •³ $f^{-1}(x) = \frac{2}{x-3}$ 	3
		<p>Method 2</p> <ul style="list-style-type: none"> •¹ write as $y = f(x)$ and start to rearrange •² express x in terms of y •³ state inverse function 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ $y = f(x) \Rightarrow x = f^{-1}(y)$ $y - 3 = \frac{2}{x}$ •² $x = \frac{2}{y-3}$ •³ $f^{-1}(y) = \frac{2}{y-3}$ $\Rightarrow f^{-1}(x) = \frac{2}{x-3}$ 	

Notes:

- In Method, 1 accept $x = \frac{2}{f^{-1}(x)} + 3$ for •¹ and •².
- In Method 2, accept ' $y - 3 = \frac{2}{x}$ ', without reference to $y = f(x) \Rightarrow x = f^{-1}(y)$ at •¹.
- In Method 2, accept $f^{-1}(x) = \frac{2}{x-3}$ without reference to $f^{-1}(y)$ at •³.
- In Method 2, beware of candidates with working where each line is not mathematically equivalent - see Candidates A and B for example.
- At •³ stage, accept f^{-1} written in terms of any dummy variable eg $f^{-1}(y) = \frac{2}{y-3}$.
- $y = \frac{2}{x-3}$ does not gain •³.
- $f^{-1}(x) = \frac{2}{x-3}$ with no working gains 3/3.
- In Method 2, where candidates make multiple algebraic errors at the •² stage, •³ is still available.

Question	Generic scheme	Illustrative scheme	Max mark
6.	(continued)		
Commonly Observed Responses:			
<p>Candidate A</p> $f(x) = \frac{2}{x} + 3$ $y = \frac{2}{x} + 3$ $y - 3 = \frac{2}{x}$ $x = \frac{2}{y - 3}$ $y = \frac{2}{x - 3}$ $f^{-1}(x) = \frac{2}{x - 3}$	<p>Candidate B</p> $f(x) = \frac{2}{x} + 3$ $y = \frac{2}{x} + 3$ $x = \frac{2}{y} + 3$ $x - 3 = \frac{2}{y}$ $y = \frac{2}{x - 3}$ $f^{-1}(x) = \frac{2}{x - 3}$	<p>•¹ ✗</p> <p>•² ✓<input type="checkbox"/></p> <p>•³ ✓<input type="checkbox"/></p>	
<p>Candidate C - BEWARE</p> $f'(x) = \dots$	<p>Candidate D</p> $x \rightarrow \frac{1}{x} \rightarrow \frac{2}{x} \rightarrow \frac{2}{x} + 3 = f(x)$ $\times 2 \rightarrow +3$ $\therefore -3 \rightarrow \div 2$ $\frac{2}{x-3} \text{ (invert)}$ $f^{-1}(x) = \frac{2}{x-3}$	<p>•³ ✗</p> <p>•¹ ✓</p> <p>•² ✓</p> <p>•³ ✓</p>	
<p>Candidate E</p> $f^{-1}(x) = \left(\frac{x-3}{2}\right)^{-1}$	<p>Candidate F</p> $f^{-1}(x) = \sqrt[2]{\frac{x-3}{2}}$	<p>•¹ ✓ •² ✓</p> <p>•³ ✓</p> <p>•¹ ✓ •² ✓</p> <p>•³ ✓</p>	
<p>Candidate G</p> $y = \frac{2}{x} + 3$ $xy = 5$ $x = \frac{5}{y}$ $f^{-1}(x) = \frac{5}{x}$ <p>However</p> $f^{-1}(x) = \frac{2+3}{x}$		<p>•¹ ✗</p> <p>•² ✓<input type="checkbox"/></p> <p>•³ ✓<input type="checkbox"/></p> <p>•³ ^</p>	

Question	Generic scheme	Illustrative scheme	Max mark
7.	<ul style="list-style-type: none"> •¹ use double angle formula to express equation in terms of $\sin x^\circ$ •² arrange in standard quadratic form •³ factorise or use quadratic formula •⁴ solve for $\sin x^\circ$ •⁵ solve for x 	<ul style="list-style-type: none"> •¹ $\dots = 3(1 - 2\sin^2 x^\circ)$ •² $6\sin^2 x^\circ + \sin x^\circ - 1 = 0$ •³ $(3\sin x^\circ - 1)(2\sin x^\circ + 1) (= 0)$ or $\sin x^\circ = \frac{-1 \pm \sqrt{25}}{12}$ •⁴ $\sin x^\circ = \frac{1}{3}$, •⁵ $\sin x^\circ = -\frac{1}{2}$ •⁵ 19.47 ..., 160.52 ..., 210, 330 	5

Notes:

1. Substituting $1 - 2\sin^2 A$ or $1 - 2\sin^2 \alpha$ for $\cos 2x^\circ$ at the •¹ stage should be treated as bad form provided the equation is written in terms of x at •² stage. Otherwise, •¹ is not available.
2. Do not penalise the omission of degree signs.
3. ‘= 0’ must appear by •³ stage for •² to be awarded. However, for candidates using the quadratic formula to solve the equation, ‘= 0’ must appear at •² stage for •² to be awarded.
4. Candidates may express the equation obtained at •² in the form $6S^2 + S - 1 = 0$, $6x^2 + x - 1 = 0$ or using any other dummy variable at the •³ stage. In these cases, award •³ for $(3S - 1)(2S + 1)$ or $(3x - 1)(2x + 1)$.
However, •⁴ is only available if $\sin x^\circ$ appears explicitly at this stage - see Candidate A.
5. The equation $1 - 6\sin^2 x^\circ - \sin x^\circ = 0$ does not gain •² unless •³ has been awarded.
6. •³ is awarded for identifying the factors of the quadratic obtained at •² eg “ $3\sin x^\circ - 1 = 0$ and $2\sin x^\circ + 1 = 0$ ”.
7. •⁴ and •⁵ are only available as a consequence of trying to solve a quadratic equation - see Candidate B.
8. •³, •⁴ and •⁵ are not available for any attempt to solve a quadratic equation written in the form $ax^2 + bx = c$ - see Candidate C.
9. •⁵ is only available where at least one of the equations at •⁴ leads to two solutions for x .
10. Do not penalise additional (correct) solutions at •⁵. However see Candidates E and F.
11. Accept answers which round to 19, 19.5 and 161.

Question	Generic scheme	Illustrative scheme	Max mark
7.	(continued)		
Commonly Observed Responses:			
Candidate A ∴ $6S^2 + S - 1 = 0$ $(3S - 1)(2S + 1) = 0$ $S = \frac{1}{3}, S = -\frac{1}{2}$ $x = 19.5, 160.5, 210, 330$	• ¹ ✓ • ² ✓ • ³ ✓ • ⁴ ^ • ⁵ <input checked="" type="checkbox"/>	Candidate B - not solving a quadratic ∴ $6 \sin^2 x^\circ + \sin x^\circ - 1 = 0$ $7 \sin x^\circ - 1 = 0$ $\sin x^\circ = \frac{1}{7}$ $x = 8.2$	• ¹ ✓ • ² ✓ • ³ ✗ • ⁴ <input checked="" type="checkbox"/> ₂ • ⁵ <input checked="" type="checkbox"/> ₂
Candidate C - not in standard quadratic form $\sin x^\circ + 2 = 3 - 6 \sin^2 x^\circ$ $6 \sin^2 x^\circ + \sin x^\circ = 1$ $\sin x^\circ (6 \sin x^\circ + 1) = 1$ $\sin x^\circ = 1 \quad 6 \sin x^\circ + 5 = 1$ $\Rightarrow \sin x = -\frac{4}{6}$ $90, 221.8, 318.2$	• ¹ ✓ • ² <input checked="" type="checkbox"/> ₂ • ³ <input checked="" type="checkbox"/> ₂ • ⁴ ✗ • ⁵ ✗	Candidate D - reading $\cos 2x^\circ$ as $\cos^2 x^\circ$ $\sin x^\circ + 2 = 3 \cos^2 x^\circ$ $\sin x^\circ + 2 = 3(1 - \sin^2 x^\circ)$ $3 \sin^2 x^\circ + \sin x^\circ - 1 = 0$ $\sin x^\circ = \frac{-1 \pm \sqrt{13}}{6}$ $\sin x^\circ = 0.434\dots, \sin x^\circ = -0.767\dots$ $25.7, 154.3, 230.1, 309.9$	• ¹ ✗ • ² <input checked="" type="checkbox"/> ₁ • ³ <input checked="" type="checkbox"/> ₁ • ⁴ <input checked="" type="checkbox"/> ₁ • ⁵ <input checked="" type="checkbox"/> ₁
Candidate E ∴ $(3 \sin x^\circ - 1)(2 \sin x^\circ + 1) = 0$ $\sin x^\circ = \frac{1}{3}, \sin x^\circ = -\frac{1}{2}$ $x = 19, x = 161 \quad x = 30$ $x = 210, x = 330$ • ⁵ ✗ However, where the final solution(s) are clearly identified by the candidate award •⁵	• ¹ ✓ • ² ✓ • ³ ✓ • ⁴ ✓ • ⁵ ✗	Candidate F ∴ $(3 \sin x^\circ - 1)(2 \sin x^\circ + 1) = 0$ $\sin x^\circ = \frac{1}{3}, \sin x^\circ = -\frac{1}{2}$ $x = 19, 161, 30, 210, 330$	• ¹ ✓ • ² ✓ • ³ ✓ • ⁴ ✓ • ⁵ ✗

Question		Generic scheme	Illustrative scheme	Max mark
8.		<p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ integrate using “upper - lower” •² identify limits •³ integrate •⁴ substitute limits •⁵ calculate shaded area 	<p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ $\int \left((x^3 - 2x^2 - 4x + 1) - (x - 5) \right) dx$ •² $\int_{-2}^1 \left((x^3 - 2x^2 - 4x + 1) - (x - 5) \right) dx$ •³ $\frac{x^4}{4} - \frac{2x^3}{3} - \frac{5x^2}{2} + 6x$ •⁴ $\left(\frac{(1)^4}{4} - \frac{2(1)^3}{3} - \frac{5(1)^2}{2} + 6(1) \right) - \left(\frac{(-2)^4}{4} - \frac{2(-2)^3}{3} - \frac{5(-2)^2}{2} + 6(-2) \right)$ •⁵ $\frac{63}{4}$ or $15\frac{3}{4}$ 	5
		<p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ know to integrate between appropriate limits for both integrals •² integrate both functions •³ substitute limits into both expressions •⁴ evaluate both integrals •⁵ evidence of subtracting areas 	<p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ $\int_{-2}^1 \dots dx$ and $\int_{-2}^1 \dots dx$ •² $\frac{x^4}{4} - \frac{2x^3}{3} - \frac{4x^2}{2} + x$ and $\frac{x^2}{2} - 5x$ •³ $\left(\frac{(1)^4}{4} - \frac{2(1)^3}{3} - \frac{4(1)^2}{2} + (1) \right) - \left(\frac{(-2)^4}{4} - \frac{2(-2)^3}{3} - \frac{4(-2)^2}{2} + (-2) \right)$ and $\left(\frac{(1)^2}{2} - 5(1) \right) - \left(\frac{(-2)^2}{2} - 5(-2) \right)$ •⁴ $-\frac{3}{4}$ and $-\frac{33}{2}$ •⁵ $-\frac{3}{4} - \left(-\frac{33}{2} \right) = \frac{63}{4}$ 	5

Question	Generic scheme	Illustrative scheme	Max mark
8.	(continued)		
Notes:			
<ol style="list-style-type: none"> 1. Correct answer with no working - award 1/5. 2. In Method 1, treat the absence of brackets at \bullet^1 stage as bad form only if the correct integral is obtained at \bullet^3 - see Candidates A and B. 3. Do not penalise lack of 'dx' at \bullet^1. 4. Limits and 'dx' must appear by the \bullet^2 stage for \bullet^2 to be awarded in Method 1 and by the \bullet^1 stage for \bullet^1 to be awarded in Method 2. 5. Where a candidate differentiates one or more terms at \bullet^3, then \bullet^3, \bullet^4 and \bullet^5 are unavailable. 6. Accept unsimplified expressions at \bullet^3 e.g. $\frac{x^4}{4} - \frac{2x^3}{3} - \frac{4x^2}{2} + x - \frac{x^2}{2} + 5x$. 7. Do not penalise the inclusion of '+c'. 8. Do not penalise the continued appearance of the integral sign after \bullet^2 9. Candidates who substitute limits without integrating do not gain \bullet^3, \bullet^4 or \bullet^5. 10. \bullet^5 is not available where solutions include statements such as '$-\frac{63}{4} = \frac{63}{4}$ square units' - see Candidate B. 11. Where a candidate only integrates $x^3 - 2x^2 - 4x + 1$ or another cubic or quartic expression, only \bullet^3 and \bullet^4 are available (from Method 1). 			

Question	Generic scheme	Illustrative scheme	Max mark
8. (continued)			
Commonly Observed Responses:			
<p>Candidate A - bad form corrected</p> $\int_{-2}^1 x^3 - 2x^2 - 4x + 1 - x - 5 dx \quad \bullet^2 \checkmark$ $= \frac{x^4}{4} - \frac{2x^3}{3} - \frac{5x^2}{2} + 6x \quad \bullet^3 \checkmark \Rightarrow \bullet^1 \checkmark$ <p>Bad form at \bullet^1 must be corrected by the integration stage and may also take the form of a missing minus sign</p>	<p>Candidate B</p> $\int_{-2}^1 x^3 - 2x^2 - 4x + 1 - x - 5 dx \quad \bullet^1 \times \bullet^2 \checkmark$ $= \frac{x^4}{4} - \frac{2x^3}{3} - \frac{5x^2}{2} - 4x \quad \bullet^3 \checkmark$ $= \left(\frac{(1)^4}{4} - \frac{2(1)^3}{3} - \frac{5(1)^2}{2} - 4(1) \right)$ $- \left(\frac{(-2)^4}{4} - \frac{2(-2)^3}{3} - \frac{5(-2)^2}{2} - 4(-2) \right) \quad \bullet^4 \checkmark$ $- \frac{57}{4} \text{ cannot be negative so } = \frac{57}{4} \quad \bullet^5 \times$ <p>However, $\int \dots = -\frac{57}{4}$ so Area = $\frac{57}{4} \quad \bullet^5 \checkmark$</p>		
<p>Candidate C - lower – upper</p> $\int_{-2}^1 \left((x-5) - (x^3 - 2x^2 - 4x + 1) \right) dx \quad \bullet^2 \checkmark$ $= \frac{x^4}{4} + \frac{2x^3}{3} + \frac{5x^2}{2} - 6x \quad \bullet^3 \checkmark$ $\left(\frac{(1)^4}{4} + \frac{2(1)^3}{3} + \frac{5(1)^2}{2} - 6(1) \right) -$ $\left(\frac{(-2)^4}{4} + \frac{2(-2)^3}{3} + \frac{5(-2)^2}{2} - 6(-2) \right) \quad \bullet^4 \checkmark$ $= \frac{63}{4}$ <p>So Area = $\frac{63}{4} \quad \bullet^1 \checkmark \bullet^5 \checkmark$</p>	<p>Candidate D - reversed limits</p> $\int_1^{-2} \left((x^3 - 2x^2 - 4x + 1) - (x-5) \right) dx \quad \bullet^1 \checkmark$ $= \frac{x^4}{4} - \frac{2x^3}{3} - \frac{5x^2}{2} + 6x \quad \bullet^3 \checkmark$ $\left(\frac{(-2)^4}{4} - \frac{2(-2)^3}{3} - \frac{5(-2)^2}{2} + 6(-2) \right)$ $- \left(\frac{(1)^4}{4} - \frac{2(1)^3}{3} - \frac{5(1)^2}{2} + 6(1) \right) \quad \bullet^4 \checkmark$ $= \frac{63}{4}$ <p>So Area = $\frac{63}{4} \quad \bullet^2 \checkmark \bullet^5 \checkmark$</p> <p>$\checkmark$</p>		
<p>Candidate E - ‘upper’ – ‘lower’</p> $= x^3 - 2x^2 - 5x + 6$ $\int_{-2}^1 (x^3 - 2x^2 - 5x + 6) dx \quad \bullet^1 \checkmark \bullet^2 \checkmark$ $= \frac{x^4}{4} - \frac{2x^3}{3} - \frac{5x^2}{2} + 6x \quad \bullet^3 \checkmark$ $\frac{37}{12} - \left(-\frac{38}{3} \right) \quad \bullet^4 \checkmark$ $= \frac{63}{4} \quad \bullet^5 \checkmark$			

Question		Generic scheme	Illustrative scheme	Max mark
9.	(a)	<ul style="list-style-type: none"> •¹ use compound angle formula •² compare coefficients •³ process for k •⁴ process for a and express in required form 	<ul style="list-style-type: none"> •¹ $k \sin x^\circ \cos a^\circ + k \cos x^\circ \sin a^\circ$ stated explicitly •² $k \cos a^\circ = -3, k \sin a^\circ = 7$ stated explicitly •³ $\sqrt{58}$ •⁴ $\sqrt{58} \sin(x + 113.19\dots)^\circ$. 	4

Notes:

1. Do not penalise the omission of degree symbols in this question.
2. Accept $k(\sin x^\circ \cos a^\circ + \cos x^\circ \sin a^\circ)$ at •¹.
3. Treat $k \sin x^\circ \cos a^\circ + \cos x^\circ \sin a^\circ$ as bad form only if the equations at the •² stage both contain k .
4. $\sqrt{58} \sin x^\circ \cos a^\circ + \sqrt{58} \cos x^\circ \sin a^\circ$ or $\sqrt{58}(\sin x^\circ \cos a^\circ + \cos x^\circ \sin a^\circ)$ are acceptable for •¹ and •³.
5. •² is not available for $k \cos x^\circ = -3$ and $k \sin x^\circ = 7$, however •⁴ may still be gained - see Candidate E.
6. •³ is only available for a single value of $k, k > 0$.
7. •⁴ is not available for a value of a given in radians.
8. Accept values of a which round to 113.
9. Candidates may use any form of the wave function for •¹, •² and •³. However, •⁴ is only available if the wave is interpreted in the form $k \sin(x + a)^\circ$.
10. Evidence for •⁴ may appear in part (b).

Question	Generic scheme	Illustrative scheme	Max mark
9. (continued)			
Commonly Observed Responses:			
<p>Candidate A</p> <p>•¹ ^</p> <p>$\sqrt{58} \cos a^\circ = -3$</p> <p>$\sqrt{58} \sin a^\circ = 7$ •² ✓ •³ ✓</p> <p>$\tan a^\circ = -\frac{7}{3}$</p> <p>$a = 113.19\dots$</p> <p>$\sqrt{58} \sin(x + 113.19\dots)^\circ$ •⁴ ✓</p>	<p>Candidate B</p> <p>$k \sin x^\circ \cos a^\circ + k \cos x^\circ \sin a^\circ$</p> <p>•¹ ✓</p> <p>$\cos a^\circ = -3$</p> <p>$\sin a^\circ = 7$ •²</p> <p>✗</p> <p>$\tan a^\circ = -\frac{7}{3}$</p> <p>$a = 113.19\dots$</p> <p>$\sqrt{58} \sin(x + 113.19\dots)^\circ$ •³ ✓ •⁴ ✗</p> <p>Not consistent with equations at •².</p>	<p>Candidate C</p> <p>$\sin x^\circ \cos a^\circ + \cos x^\circ \sin a^\circ$ •¹ ✗</p> <p>$\cos a^\circ = -3$</p> <p>$\sin a^\circ = 7$ •² <input checked="" type="checkbox"/></p> <p>$k = \sqrt{58}$ •³ ✓</p> <p>$\tan a^\circ = -\frac{7}{3}$</p> <p>$a = 113.19\dots$</p> <p>$\sqrt{58} \sin(x + 113.19\dots)^\circ$ •⁴ ✗</p>	
<p>Candidate D - errors at •²</p> <p>$k \sin x \cos a + k \cos x \sin a$ •¹ ✓</p> <p>$k \cos a^\circ = 7$</p> <p>$k \sin a^\circ = -3$ •² ✗</p> <p>$\tan a^\circ = -\frac{3}{7}$</p> <p>$a = 336.80\dots$</p> <p>$\sqrt{58} \sin(x + 336.80\dots)^\circ$ •³ ✓ •⁴ <input checked="" type="checkbox"/></p>	<p>Candidate E - use of x at •²</p> <p>$k \sin x \cos a + k \cos x \sin a$ •¹ ✓</p> <p>$k \cos x^\circ = -3$</p> <p>$k \sin x^\circ = 7$ •² ✗</p> <p>$\tan a^\circ = -\frac{7}{3}$</p> <p>$a = 113.19\dots$</p> <p>$\sqrt{58} \sin(x + 113.19\dots)^\circ$</p> <p>•³ ✓ •⁴ <input checked="" type="checkbox"/></p>	<p>Candidate F</p> <p>$k \sin A \cos B + k \cos A \sin B$ •¹ ✗</p> <p>$k \cos A = -3$</p> <p>$k \sin A = 7$ •² ✗</p> <p>$\tan A = -\frac{7}{3}$</p> <p>$A = 113.19\dots$</p> <p>$\sqrt{58} \sin(x + 113.19\dots)^\circ$ •³ ✓ •⁴ <input checked="" type="checkbox"/></p>	

Question			Generic scheme	Illustrative scheme	Max mark
9.	(b)	(i)	• ⁵ state maximum value	• ⁵ $2\sqrt{58}$	1
		(ii)	<p style="text-align: center;">Method 1</p> <p>•⁶ start to solve</p> <p>•⁷ state value of x</p> <p style="text-align: center;">Method 2</p> <p>•⁶ start to solve</p> <p>•⁷ state value of x</p>	<p style="text-align: center;">Method 1</p> <p>•⁶ $x + 113.19\dots = 90$ leading to $x = -23.19\dots$</p> <p>•⁷ $x = 336.80\dots$</p> <p style="text-align: center;">Method 2</p> <p>•⁶ $x + 113.19\dots = 450$</p> <p>•⁷ $x = 336.80\dots$</p>	2
Notes:					
11. • ⁷ is only available where an angle outwith the range $0 \leq x < 360$ needs to be considered - see Candidate G.					
12. • ⁷ is only available where • ⁶ has been awarded. However, see Candidate K.					
Commonly Observed Responses:					
Candidate G - not considering angle outwith $0 \leq x < 360$			Candidate H - simplification		
$\sqrt{58} \sin(x - 23)^\circ$ from part (a)			(i) $2\sqrt{58}$ •⁵ ✓		
$x - 23 = 90$ •⁶ <input checked="" type="checkbox"/> 1			(ii) $\sqrt{58} \sin(x + 113)^\circ = \sqrt{58}$		
$x = 113$ •⁷ <input checked="" type="checkbox"/> 2			$x + 113 = 90$		
			$x = -23$ •⁶ ✓		
			$x = 337$ •⁷ ✓		
Candidate I - follow-through marking			Candidate J - graphical approach		
(i) $\sqrt{58}$ •⁵ ✗			(i) $\sqrt{58}$ •⁵ ✗		
(ii) $2\sqrt{58} \sin(x + 113)^\circ = \sqrt{58}$			(ii) max occurs when $x + 113 = 90$		
$x + 113 = 30$			$x = -23$ •⁶ ✓		
$x = -83$ •⁶ <input checked="" type="checkbox"/> 1			$x = 337$ •⁷ ✓		
$x = 277$ •⁷ <input checked="" type="checkbox"/> 1					
Candidate K - no acknowledgement of $\times 2$					
(i) $\sqrt{58}$ •⁵ ✗					
(ii) $\sqrt{58} \sin(x + 113)^\circ = \sqrt{58}$					
$x + 113 = 90$					
$x = -23$ •⁶ ✗					
$x = 337$ •⁷ <input checked="" type="checkbox"/> 1					

Question		Generic scheme	Illustrative scheme	Max mark											
10.		<p>Method 1</p> <ul style="list-style-type: none"> •¹ differentiate one term •² complete differentiation and interpret condition •³ determine zeros of quadratic expression •⁴ state range with justification 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $6x^2 \dots$ or $+18x \dots$ or -24 •² $6x^2 + 18x - 24 < 0$ •³ 1 and -4 •⁴ $-4 < x < 1$ with eg labelled sketch 	4											
		<p>Method 2</p> <ul style="list-style-type: none"> •¹ differentiate one term •² complete differentiation and determine zeros of quadratic expression •³ construct nature table(s) •⁴ interpret sign of derivative and state range 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ $6x^2 \dots$ or $+18x \dots$ or -24 •² $6x^2 + 18x - 24$ and 1 and -4 •³ <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>\dots</td> <td>-4</td> <td>\dots</td> <td>1</td> <td>\dots</td> </tr> <tr> <td>$f'(x)$</td> <td>$+$</td> <td>0</td> <td>$-$</td> <td>0</td> <td>$+$</td> </tr> </table> •⁴ decreasing when $f'(x) < 0$ so $-4 < x < 1$ 	x	\dots	-4	\dots	1	\dots	$f'(x)$	$+$	0	$-$	0	$+$
x	\dots	-4	\dots	1	\dots										
$f'(x)$	$+$	0	$-$	0	$+$										

Notes:

1. At •³ do not penalise candidates who fail to extract the common factor or who have divided the quadratic inequality by 6.
2. •³ and •⁴ are not available to candidates who arrive at a linear expression at •².
3. Accept the appearance of -4 and 1 within inequalities for •³.
4. At •⁴, accept " $x > -4, x < 1$ " together with the required justification.

Commonly Observed Responses:

<p>Candidate A</p> $6x^2 + 18x - 24 < 0$ • ¹ ✓ • ² ✓ $6x^2 + 18x - 24 = 0$ • ³ ✓ $x = -4, 1$ • ⁴ ✓ $-4 < x < 1$ with sketch	<p>Candidate B - no initial inequation</p> $6x^2 + 18x - 24 = 0$ • ¹ ✓ • ² ✗ $x = -4, 1$ • ³ ✓ $-4 < x < 1$ with sketch • ⁴ ✗
<p>Candidate C</p> Decreasing when $f'(x) < 0$ $f'(x) = 6x^2 + 18x - 24$ • ¹ ✓ • ² ✓ \vdots	<p>Candidate D - condition applied after simplification</p> $f'(x) = 6x^2 + 18x - 24$ • ¹ ✓ $x^2 + 3x - 4 < 0$ • ² ^ $x = -4, 1$ • ³ ✓ $-4 < x < 1$ with sketch • ⁴ ✓

Question		Generic scheme	Illustrative scheme	Max mark
11.	(a)	<ul style="list-style-type: none"> •¹ state centre of C_1 •² state centre of C_2 •³ calculate distance between centres 	<ul style="list-style-type: none"> •¹ $(4, -2)$ •² $(-1, 3)$ •³ $\sqrt{50}$ or $5\sqrt{2}$ or 7.07... 	3
Notes:				
1. Accept $x = 4, y = -2$ for • ¹ and $x = -1, y = 3$ • ² . Do not accept $g = 1, f = -3$ for • ² . 2. Do not penalise lack of brackets in • ¹ and • ² .				
Commonly Observed Responses:				
	(b)	<ul style="list-style-type: none"> •⁴ state radius of C_1 •⁵ calculate radius of C_2 •⁶ demonstrate and communicate result 	<ul style="list-style-type: none"> •⁴ $r_1 = \sqrt{37}$ or 6.08... •⁵ $r_2 = \sqrt{17}$ or 4.12... •⁶ $10.20... > 7.07... (> 1.95...)$ \therefore circles intersect at two distinct points 	3
Notes:				
3. Accept $\sqrt{1^2 + 3^2 + 7} = \sqrt{17}$ or $\sqrt{1^2 + (-3)^2 + 7} = \sqrt{17}$ for • ⁵ . However, do not accept $\sqrt{(-1)^2 + 3^2 + 7} = \sqrt{17}$. 4. At • ⁶ comparison must be made using decimals. Do not accept $\sqrt{37} + \sqrt{17} > \sqrt{50}$ without any further working. 5. Evidence for • ⁴ and • ⁵ may be found in part (a). 6. For candidates who use simultaneous equations, award • ⁴ for substitution of $y = x + 1$ into the equation of one of the circles, • ⁵ for rearranging in standard quadratic form and • ⁶ for obtaining distinct x -coordinates. 7. Do not penalise the omission of “at two distinct points” at • ⁶ .				
Commonly Observed Responses:				

Question	Generic scheme	Illustrative scheme	Max mark
12.	<ul style="list-style-type: none"> •¹ integrate one term •² complete integration •³ substitute for x and y •⁴ state expression for y 	<ul style="list-style-type: none"> •¹ eg $\frac{8x^4}{4} \dots$ •² eg $\dots + 3x + c$ •³ $3 = \frac{8 \times (-1)^4}{4} + 3 \times (-1) + c$ •⁴ $y = 2x^4 + 3x + 4$ 	4

Notes:

1. For candidates who omit $+c$ only •¹ is available.
2. For candidates who differentiate either term, •², •³, and •⁴ are not available.
3. Do not penalise the appearance of an integral sign and/or dx at •² and •³.

Commonly Observed Responses:

Candidate A - incomplete substitution

$y = 2x^4 + 3x + c$ •¹ ✓ •² ✓
 $y = 2(-1)^4 + 3(-1) + c$
 $c = 4$ •³ ^
 $y = 2x^4 + 3x + 4$ •⁴ ✓₁

Candidate B - partial integration

$y = 2x^4 + 3 + c$ •¹ ✓ •² ✗
 $3 = 2(-1)^4 + 3 + c$ •³ ✓₁
 $c = -2$
 $y = 2x^4 + 1$ •⁴ ✓₁

Candidate C - integrating over two lines

$y = 2x^4 + 3x$ •¹ ✓ •² ✗
 $y = 2x^4 + 3x + c$
 $3 = 2(-1)^4 + 3(-1) + c$ •³ ✓
 $y = 2x^4 + 3x + 4$ •⁴ ✓

Question		Generic scheme	Illustrative scheme	Max mark
13.	(a)	• ¹ calculate concentration	• ¹ 9.38... (mg/l)	1
Notes:				
1. Accept any answer which rounds to 9.4 for • ¹ .				
Commonly Observed Responses:				
	(b)	• ² substitute • ³ write in logarithmic form • ⁴ process for t	• ² $0.66 = 11 \times e^{-0.0053 t}$ • ³ $\log_e \frac{0.66}{11} = -0.0053t$ • ⁴ 530.83... (minutes)	3
Notes:				
2. Where values other than 0.66 are used in the substitution, • ³ and • ⁴ are still available. 3. Evidence for • ³ must be stated explicitly. 4. At • ³ all exponentials must be processed. 5. Any base may be used at • ³ stage - see Candidate A. 6. Accept $\ln 0.06 = -0.0053t \ln e$ for • ³ . 7. Accept any answer where $530 \leq t \leq 532$ at • ⁴ . 8. • ⁴ is unavailable where a candidate rounds the value of $\ln 0.06$ to fewer than 2 decimal places. 9. The calculation at • ⁴ must follow from the valid use of exponentials and logarithms at • ² and • ³ . 10. For candidates with no working or who take an iterative approach to arrive at $t = 532$, $t = 531$ or $t = 530$ award 1/3. However, if, in any iterations C_t is evaluated for $t = 530$ and $t = 531$ leading to a final answer of $t = 531$ (minutes) then award 3/3.				
Commonly Observed Responses:				
Candidate A		Candidate B		
$0.66 = 11e^{-0.0053t}$ • ² ✓ $0.06 = e^{-0.0053t}$ $\log_{10} 0.06 = -0.0053t \log_{10} e$ • ³ ✓ $t = 531$ minutes • ⁴ ✓		$0.66 = 11e^{-0.0053t}$ • ² ✓ $t = 531$ minutes • ³ ^ • ⁴ <input checked="" type="checkbox"/> 1		

Question			Generic scheme	Illustrative scheme	Max mark
14.	(a)	(i)	• ¹ express A in terms of x and h	• ¹ $(A =) 6x^2 + 10xh$	1
		(ii)	• ² express h in terms of x • ³ substitute for h and demonstrate result	• ² $h = \frac{7200 - 6x^2}{10x}$ • ³ $V = 3x \times 2x \times \left(\frac{7200 - 6x^2}{10x} \right)$ leading to $V = 4320x - \frac{18}{5}x^3$	2

Notes:

1. Accept unsimplified expressions for •¹.
2. •² is only available where the (simplified) expression for A contains at least 2 terms.
3. The substitution for h at •³ must be clearly shown for •³ to be awarded.

Commonly Observed Responses:

	(b)		• ⁴ differentiate • ⁵ equate expression for derivative to 0 • ⁶ solve for x • ⁷ verify nature	• ⁴ $4320 - \frac{54}{5}x^2$ • ⁵ $4320 - \frac{54}{5}x^2 = 0$ • ⁶ 20 • ⁷ table of signs for a derivative <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td style="padding: 5px;">x</td> <td style="padding: 5px;">...</td> <td style="padding: 5px;">20</td> <td style="padding: 5px;">...</td> </tr> <tr> <td style="padding: 5px;">$V'(x)$</td> <td style="padding: 5px;">+</td> <td style="padding: 5px;">0</td> <td style="padding: 5px;">-</td> </tr> <tr> <td style="padding: 5px;">shape</td> <td style="padding: 5px;">/</td> <td style="padding: 5px;">—</td> <td style="padding: 5px;">\</td> </tr> </table> \therefore maximum (when $x = 20$)	x	...	20	...	$V'(x)$	+	0	-	shape	/	—	\	4
x	...	20	...														
$V'(x)$	+	0	-														
shape	/	—	\														

Notes:

4. For any approach which does not use differentiation award 0/4.
5. •⁵ can be awarded for $\frac{54}{5}x^2 = 4320$.
6. For candidates who integrate any term at the •⁴ stage, only •⁵ is available on follow through for setting their 'derivative' to 0.
7. Ignore the appearance of -20 at mark •⁶.
8. Where -20 is considered in a nature table (or second derivative), " $x = 20$ " must be clearly identified as leading to the maximum area.
9. •⁶ and •⁷ are not available to candidates who state that the maximum exists at a negative value of x .
10. Do not penalise statements such as "max volume is 20" or "max is 20" at •⁷.

Question	Generic scheme	Illustrative scheme	Max mark																																				
14. (continued)																																							
Commonly Observed Responses:																																							
Candidate A - second derivative $V''(x) = -\frac{108}{5}x$ $V''(20) < 0 \therefore$ maximum	<p style="text-align: right;">•⁷ ✓</p>	Candidate B - beware of multiplying before equating $V'(x) = 4320 - \frac{54}{5}x^2$ $V'(x) = 21600 - 54x^2$ $21600 - 54x^2 = 0$ $x = 20$ <input checked="" type="checkbox"/> 1	<p style="text-align: right;">•⁴</p> <p style="text-align: right;">•⁵ ✗</p> <p style="text-align: right;">•⁶</p>																																				
Candidate C Stationary points when $V'(x) = 0$ $V'(x) = 4320 - \frac{54}{5}x^2$	<p style="text-align: right;">•⁴ ✓ •⁵ ✓</p>																																						
For the table of signs for a derivative, accept: <table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td>x</td> <td>20^-</td> <td>20</td> <td>20^+</td> <td>x</td> <td>\rightarrow</td> <td>20</td> <td>\rightarrow</td> <td>x</td> <td>a</td> <td>20</td> <td>b</td> </tr> <tr> <td>$V'(x)$</td> <td>+</td> <td>0</td> <td>-</td> <td>$V'(x)$</td> <td>+</td> <td>0</td> <td>-</td> <td>$V'(x)$</td> <td>+</td> <td>0</td> <td>-</td> </tr> <tr> <td>Slope or shape</td> <td></td> <td></td> <td></td> <td>Slope or shape</td> <td></td> <td></td> <td></td> <td>Slope or shape</td> <td></td> <td></td> <td></td> </tr> </table> <p style="text-align: center;">Arrows are taken to mean 'in the neighbourhood of' Where $a < 20$ and $b > 20$</p>				x	20^-	20	20^+	x	\rightarrow	20	\rightarrow	x	a	20	b	$V'(x)$	+	0	-	$V'(x)$	+	0	-	$V'(x)$	+	0	-	Slope or shape				Slope or shape				Slope or shape			
x	20^-	20	20^+	x	\rightarrow	20	\rightarrow	x	a	20	b																												
$V'(x)$	+	0	-	$V'(x)$	+	0	-	$V'(x)$	+	0	-																												
Slope or shape				Slope or shape				Slope or shape																															
For the table of signs for a derivative, accept: <table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td>x</td> <td>\rightarrow</td> <td>-20</td> <td>\rightarrow</td> <td>20</td> <td>\rightarrow</td> <td>x</td> <td>a</td> <td>-20</td> <td>b</td> <td>20</td> <td>c</td> </tr> <tr> <td>$V'(x)$</td> <td>-</td> <td>0</td> <td>+</td> <td>0</td> <td>-</td> <td>$V'(x)$</td> <td>-</td> <td>0</td> <td>+</td> <td>0</td> <td>-</td> </tr> <tr> <td>Slope or shape</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>Slope or shape</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </table> <p style="text-align: center;">Since the function is continuous $-20 \rightarrow 20$ is acceptable Since the function is continuous $-20 < b < 20$ is acceptable</p>				x	\rightarrow	-20	\rightarrow	20	\rightarrow	x	a	-20	b	20	c	$V'(x)$	-	0	+	0	-	$V'(x)$	-	0	+	0	-	Slope or shape						Slope or shape					
x	\rightarrow	-20	\rightarrow	20	\rightarrow	x	a	-20	b	20	c																												
$V'(x)$	-	0	+	0	-	$V'(x)$	-	0	+	0	-																												
Slope or shape						Slope or shape																																	
<ul style="list-style-type: none"> For this question do not penalise the omission of 'x' or the word 'shape'/'slope'. Stating values of $V'(x)$ is an acceptable alternative to writing '+' or '-' signs. Acceptable variations of $V'(x)$ are: V', $\frac{dV}{dx}$, and $4320 - \frac{54}{5}x^2$. Accept $\frac{dy}{dx}$ only where candidates have previously used $y = 4320x - \frac{18}{5}x^3$ in their working. 																																							

Question	Generic scheme	Illustrative scheme	Max mark	
15.	<ul style="list-style-type: none"> •¹ determine gradient of tangent •² determine gradient of radius •³ strategy to find centre •⁴ state coordinates of centre 	<ul style="list-style-type: none"> •¹ $-\frac{1}{3}$ •² 3 •³ eg $y = 3x - 1$ or $3 = \frac{y-5}{x-2}$ •⁴ (0, -1) 	4	
Notes:				
<ol style="list-style-type: none"> 1. Ignore errors in processing the constant term in •¹. 2. Do not accept $m = -\frac{1}{3}x$ for •¹. However •², •³ and •⁴ are still available where the candidate uses a numerical value for m_{\perp}. 3. Accept $y-5 = 3(x-2)$ as evidence for •³. 4. •⁴ is only available as a consequence of trying to find and use a perpendicular gradient along with a point on the y-axis. 5. Where candidates use “stepping out” with the perpendicular gradient, the diagram must be consistent with the solution to gain •³ and •⁴. 6. Accept “$x = 0$”, “$y = -1$” stated explicitly for •⁴. 				
Commonly Observed Responses:				
<p>Candidate A - perpendicular gradient clearly stated</p> $x + 3y = 17$	$m_{\perp} = 3$ $y = 3x - 1$	<ul style="list-style-type: none"> •¹ ✓ •² ✓ •³ ✓ 	<p>Candidate B - no communication for perpendicular gradient</p> $x + 3y = 17$ $y = -\frac{1}{3}x + \frac{17}{3}$ $m = 3$ $y = 3x - 1$	<ul style="list-style-type: none"> •¹ ^ •² ^ •³ ✓₁ •⁴ is available
<p>Candidate C - no communication for perpendicular gradient or rearrangement</p> $x + 3y = 17$ $m = 3$ $y = 3x - 1$	<ul style="list-style-type: none"> •¹ ^ •² ^ •³ ✓₂ •⁴ is not available 	<p>Candidate D - using geometry</p> <p>∴ Using point diametrically opposite (2,5), by symmetry identify that x-coordinate is -2.</p> <p>∴ $y = 3(-2) - 1 = -7$.</p> <p>Centre is midpoint of (-2, -7) and (2,5).</p> <p>∴ centre is (0, -1)</p>	<ul style="list-style-type: none"> •¹ ✓ •² ✓ •³ ✓ •⁴ ✓ 	
<p>Candidate E - incorrect gradient</p> $x + 3y = 17$ $3y = -x + 17$ $m_{\perp} = 1$ $1 = \frac{5-y}{2-0}$ Centre is at (0,3)	<ul style="list-style-type: none"> •¹ ^ •² x •³ ✓₁ •⁴ ✓₁ 			

[END OF MARKING INSTRUCTIONS]