		FOR OFFICIAL USE National Qualificati 2015	ons						Mark		
X757/76/ TUESDAY, 5 MAY 1:00 PM – 3:30 P/	01 w				Sect	ion 1	1 – a IIII *	Ansynd S		nysics r Gric tion 2	s 1 2 ₩
Fill in these bo	xes and reac	l what is print	ed belo	ow.	Town						
Forename(s)	Forename(s) Surname							Num	nber o	of seat	
Date of birt Day Total marks — 1	Month	Year	Scott	tish cai	ndidat	e numbe	er				
SECTION 1 — 20 Attempt ALL que Instructions for SECTION 2 — 1 Attempt ALL que) marks estions. the completi 10 marks estions.	ion of Section	are gi	ven on	Page :	two.			757/7	// /02	
Reference may to the Relations Care should be calculations. Write your answ and rough work identify the qu booklet. You sh Use blue or blac Before leaving t Invigilator; if yo	be made to hip Sheet X7 taken to give rers clearly in is provided estion numb ould score th ck ink. he examinat ou do not, yo	the Data Sheef 57/76/11. In the spaces pl at the end of er you are at mough your rou ion room you n u may lose all t	e numb rovided this bo cemptir igh wor nust giv the mai	ge two per of s in this oklet. ng. Ar k when k when re this rks for	signific s book If you ny rou ny ou h bookle this pa	e questi ant figu let. Ado u use th gh work nave wri at to the aper.	on pa res ir ditior is spa mus tten	aper X In the fi nal space ace you st be v your fi	nal an ce for u mu writte nal co	r answers t r answers st clearl en in thi opy.	a rs y is
L		* X 7 5	776	0 1 0) 1 *						©

The questions for Section 1 are contained in the question paper X757/76/02. Read these and record your answers on the answer grid on *Page three* opposite. Use **blue** or **black** ink. Do NOT use gel pens or pencil.

- 1. The answer to each question is **either** A, B, C, D or E. Decide what your answer is, then fill in the appropriate bubble (see sample question below).
- 2. There is only one correct answer to each question.
- 3. Any rough work must be written in the additional space for answers and rough work at the end of this booklet.

Sample Question

The energy unit measured by the electricity meter in your home is the:

- A ampere
- B kilowatt-hour
- C watt
- D coulomb
- E volt.

The correct answer is B-kilowatt-hour. The answer B bubble has been clearly filled in (see below).



Changing an answer

If you decide to change your answer, cancel your first answer by putting a cross through it (see below) and fill in the answer you want. The answer below has been changed to **D**.



If you then decide to change back to an answer you have already scored out, put a tick (\checkmark) to the **right** of the answer you want, as shown below:







С D Ε Α В \bigcirc \bigcirc Ο \bigcirc \bigcirc 1 2 Ο Ο Ο Ο Ο \bigcirc Ο \bigcirc 3 \bigcirc \bigcirc Ο Ο Ο 4 Ο Ο \bigcirc 5 \bigcirc \bigcirc \bigcirc \bigcirc \bigcirc Ο Ο Ο Ο 6 \bigcirc 7 Ο Ο Ο Ο 8 Ο Ο Ο Ο Ο 9 Ο \bigcirc \bigcirc \bigcirc \bigcirc 10 Ο Ο Ο Ο Ο Ο Ο Ο 11 \bigcirc Ο Ο Ο Ο Ο Ο 12 \bigcirc Ο \bigcirc 13 \bigcirc Ο Ο Ο Ο Ο Ο 14 15 \bigcirc \bigcirc Ο \bigcirc \bigcirc Ο 16 Ο Ο Ο Ο 17 Ο Ο \bigcirc Ο Ο Ο Ο Ο Ο Ο 18 19 \bigcirc \bigcirc Ο \bigcirc \bigcirc 20 Ο Ο Ο Ο Ο



[BLANK PAGE]

DO NOT WRITE ON THIS PAGE



L

[Turn over for SECTION 2 on Page six

DO NOT WRITE ON THIS PAGE











MARKS DO NOT THIS A student sets up an experiment to investigate collisions between two 2. trolleys on a long, horizontal track. force laptop sensor 1 • 2 m s^{−1} lightgate lightgate $0.60 \,\mathrm{m\,s^{-1}}$ trolley X mask trolley Y The mass of trolley X is 0.25 kg and the mass of trolley Y is 0.45 kg. The effects of friction are negligible. In one experiment, trolley X is moving at 1.2 m s^{-1} to the right and trolley Y is moving at 0.60 m s^{-1} to the left. The trolleys collide and do not stick together. After the collision, trolley X rebounds with a velocity of $0.80 \,\mathrm{m\,s^{-1}}$ to the left. (a) Determine the velocity of trolley Y after the collision. 3 Space for working and answer [Turn over





2. (b) (continued)

(iii) Sketch a velocity-time graph to show how the velocity of trolley X varies from 0.50 s before the collision to 0.50 s after the collision.

Numerical values are required on both axes. You may wish to use the square-ruled paper on *Page thirty-six*.

[Turn over

MARKS DO NOT WRITE IN THIS MARGIN





[Turn over for Question 4 on Page fourteen

DO NOT WRITE ON THIS PAGE





Г			
4. (b) (continued)		MARKS	DO NOT WRITE IN THIS MARGIN
(ii) Show that the redshift of the light from the distant 0.098.	galaxy	is 2	
Space for working and answer			
(iii) Calculate the approximate distance to the distant galaxy	•	5	
Space for working and answer			
1	Turn ov	er	

5.	A quote from a well-known science fiction writer states:	MARKS	DO NOT WRITE IN THIS MARGIN
	"In the beginning there was nothing, which exploded."		
	Using your knowledge of physics, comment on the above statement.	3	





MARKS DO NOT WRITE IN THIS MARGIN

7. The use of analogies from everyday life can help better understanding of physics concepts. Throwing different balls at a coconut shy to dislodge a coconut is an analogy which can help understanding of the photoelectric effect.



Use your knowledge of physics to comment on this analogy.



[Turn over for Question 8 on Page twenty

DO NOT WRITE ON THIS PAGE



- MARKS DO NOT WRITE IN THIS MARGIN
- 8. A student investigates how irradiance I varies with distance d from a point source of light.



The distance between a small lamp and a light sensor is measured with a metre stick. The irradiance is measured with a light meter.

The apparatus is set up as shown in a darkened laboratory.

The following results are obtained.

<i>d</i> (m)	0.20	0.30	0.40	0.50
$I (W m^{-2})$	134.0	60.5	33.6	21.8

- (a) State what is meant by the term *irradiance*.
- (b) Use all the data to establish the relationship between irradiance ${\cal I}$ and distance d .







MARKS DO NOT WRITE IN THIS MARGIN 9. A student carries out two experiments to investigate the spectra produced from a ray of white light. (a) In the first experiment, a ray of white light is incident on a glass prism as shown. not to scale normal 60° spectrum 42° ray of white light air glass (i) Explain why a spectrum is produced in the glass prism. 1 (ii) The refractive index of the glass for red light is 1.54. Calculate the speed of red light in the glass prism. 3 Space for working and answer





[BLANK PAGE]

DO NOT WRITE ON THIS PAGE



L



* X 7 5 7 7 6 0 1 2 5 *

10. (continued)

(b) A technician sets up the following circuit with a different car battery connected to a variable resistor R.



Readings of current I and terminal potential difference V from this circuit are used to produce the following graph.





10	(b)	(con	tinued)	Μ	ARKS	DO NOT WRITE IN THIS	
10.	Use information from the graph to determine:						
		(i)	the e.m.f. of the battery;		1		
			Space for working and answer				
			space for working and answer				
					_		
		(ii)	the internal resistance of the battery;		3		
			Space for working and answer				
				[Turn over			
L						. ,	
			* X 7 5 7 7 6 0 1 2 7 *				
	Loo	king f	or more resources? Visit <u>https://sqa.my/</u> - Scotland's #1	Past Paper Ar	chive		F

10. (b) (continued)

(iii) After being used for some time the e.m.f. of the battery decreases to 11.5V and the internal resistance increases to 0.090 Ω.

The battery is connected to a battery charger of constant e.m.f. 15.0 V and internal resistance of 0.45Ω as shown.



(A) Switch S is closed. Calculate the initial charging current. Space for working and answer

Explain why the charging current decreases as the battery (B) charges.

2

3



[Turn over for Question 11 on Page thirty

DO NOT WRITE ON THIS PAGE



11. A defibrillator is a device that provides a high energy electrical impulse to correct abnormal heart beats.



The diagram shows a simplified version of a defibrillator circuit.



The switch is set to position 1 and the capacitor charges.

(a) Show the charge on the capacitor when it is fully charged is 0.16 C.

Space for working and answer



MARKS DO NOT WRITE IN THIS MARGIN

11. (continued)

(b) Calculate the maximum energy stored by the capacitor.

Space for working and answer

(c) To provide the electrical impulse required the capacitor is discharged through the person's chest using the paddles as shown



The initial discharge current through the person is 35.0A.

(i) Calculate the effective resistance of the part of the person's body between the paddles.

Space for working and answer



MARKS DO NOT WRITE IN THIS MARGIN

3

11. (c) (continued)

(ii) The graph shows how the current between the paddles varies with time during the discharge of the capacitor.



The effective resistance of the person remains the same during this time.

Explain why the current decreases with time.

(iii) The defibrillator is used on a different person with larger effective resistance. The capacitor is again charged to 2.50 kV.

On the graph in (c)(ii) add a line to show how the current in this person varies with time.

(An additional graph, if required, can be found on *Page thirty-eight*).

2

1



MARKS DO NOT THIS 12. A student carries out an investigation to determine the refractive index of a prism. A ray of monochromatic light passes through the prism as shown. not to scale 60 Ddeviated θ ray incident ray 60° The angle of deviation D is the angle between the direction of the incident ray and the deviated ray. The student varies the angle of incidence θ and measures the corresponding angles of deviation D. The results are shown in the table. Angle of incidence θ (°) Angle of deviation D (°) 30.0 47.0 40.0 38.1 50.0 37.5 60.0 38.8 70.0 42.5 (a) Using the square-ruled paper on Page thirty-five, draw a graph of Dagainst θ . 3 (b) Using your graph state the two values of θ that produce an angle of deviation of 41.0° . 1 (c) Using your graph give an estimate of the minimum angle of deviation D_{m} . 1



12. (continued)

(d) The refractive index n of the prism can be determined using the relationship.

$$n\sin\left(\frac{A}{2}\right) = \sin\left(\frac{A+D_m}{2}\right)$$

where A is the angle at the top of the prism, and $D_{\rm m}$ is the minimum angle of deviation.

Use this relationship and your answer to (c) to determine the refractive index of the prism.

Space for working and answer

(e) Using the same apparatus, the student now wishes to determine more precisely the minimum angle of deviation.

Suggest two improvements to the experimental procedure that would achieve this.

2

[END OF QUESTION PAPER]



MARKS DO NOT WRITE IN THIS MARGIN













ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK

Additional graph for Question 11 (c)(iii)





ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK



ACKNOWLEDGEMENT

Section 2, Question 7-daseaford/shutterstock.com

